# Randić Structure of a Graph

Juan Rada and Carlos Uzcátegui Departamento de Matemáticas, Facultad de Ciencias Universidad de Los Andes, 5101 Mérida, Venezuela e-mail: juanrada@ciens.ula.ve and uzca@ciens.ula.ve

#### Abstract

Let  $\mathcal{G}$  be a collection of graphs with n vertices. We present a simple description of  $[G]_{\chi} = \{H \in \mathcal{G} : \chi(H) = \chi(G)\}$  where  $\chi$  denotes the Randić index. We associate to  $\mathcal{G}$  a  $\mathbb{Q}$ -linear map  $\rho : \mathbb{Q}^m \to \mathbb{Q}^k$  (for some integers k, m depending on  $\mathcal{G}$ ) such that the kernel of  $\rho$  contains the necessary information to describe  $[G]_{\chi}$  in terms of linear equations. These results provide precise tools for analyzing the behavior of  $\chi$  on a collection of graphs.

#### 1 Introduction

Let G be a simple graph (i.e. G does not have loops or multiple edges) with n vertices. The connectivity index of G, denoted by  $\chi$ , is defined as follows

$$\chi(G) = \sum_{1 \le i \le j \le n-1} \frac{m_{ij}(G)}{\sqrt{ij}} \tag{1}$$

where  $m_{ij}(G)$  is the number of edges in G between vertices with degrees i and j.

Randić ([14]) introduced this index (known today as the Randić index) in the study of branching properties of alkanes, and it became one of the most useful graph-based molecular descriptors in applications to physical and chemical properties ([10, 11]). In spite of this great number of practical applications, the study of the general mathematical properties of  $\chi$  started recently (see for instance [1, 2, 3, 4, 5, 6, 7, 9, 12, 13]).

It is well known that  $\chi$  does not separate non-isomorphic graphs and, of course, it does not distinguish between graphs with equal  $m_{ij}$ 's. So, given a (significant) collection of graphs  $\mathcal{G}$  with *n* vertices and  $G \in \mathcal{G}$ , it would be interesting to describe the set

$$[G]_{\chi} = \{ H \in \mathcal{G} : \chi(H) = \chi(G) \}$$

To compute all graphs in this set seems a quite demanding task, since  $[G]_{\chi}$  is quite complex from a combinatoric view point. In spite of all this, we will show that  $[G]_{\chi}$  can be described very simply in terms of a system of linear equations. In order to state our results we need the following identity for the Randić index ([8])

$$\chi(G) = \frac{n(G)}{2} - \frac{1}{2} \sum_{1 \le i < j \le n-1} \left(\frac{1}{\sqrt{j}} - \frac{1}{\sqrt{i}}\right)^2 \cdot m_{ij}(G)$$
(2)

Notice that the  $m_{ii}$ 's are not included. We denote by R(G) the set  $\{m_{ij}(G)\}_{1 \le i < j \le n-1}$ and called it the *Randić structure* of G. It is clear from (2) that if R(G) = R(G'), then  $\chi(G) = \chi(G')$ . Our description of  $[G]_{\chi}$  will be in terms of the Randić structure of a graph.

The basic idea is the following. Let  $\mathcal{G}$  be a collection of graphs and  $G \in \mathcal{G}$ . First we express  $\chi(G)$  as a linearly independent combination of certain  $\sqrt{q_i}$ 's, where the  $q_i$ 's are positive integers which derive from the set of vertex degrees of graphs from  $\mathcal{G}$ . The coefficients in this linear combination, which will depend on R(G), induces in a natural way a  $\mathbb{Q}$ -linear map  $\rho : \mathbb{Q}^m \to \mathbb{Q}^k$  (for some integers k, m depending on  $\mathcal{G}$ ). As we shall see, the kernel of  $\rho$  contains precise information that will lead us to characterize the set  $[G]_{\chi}$ , in terms of a system of linear equations on the  $m_{ij}$ 's. More precisely, we will show that for all  $G, H \in \mathcal{G}$ 

$$\chi(G) = \chi(H) \Longleftrightarrow R(G) - R(H) \in \ker \rho$$

Perhaps the simplest situation, regarding the problem of describing  $[G]_{\chi}$ , is when  $[G]_{\chi}$ consists merely of those  $H \in \mathcal{G}$  such that R(H) = R(G). In this case, we will say  $\mathcal{G}$  has *Randić* structure property (RSP). We will show a fairly general method to generate collections with the RSP. Furthermore, it will be shown that any collection of graphs  $\mathcal{G}$  can be decomposed into pairwise disjoint subcollections with the RSP. Moreover (and this is the non trivial part of such decompositions) the subcollections are defined by linear equations on the  $m_{ij}$ 's. As an example of how these ideas can be used to get information about the behavior of  $\chi$  on a collection of graphs, we will make in the last section an analysis of the collection of branch regular trees of degree 4 (i.e. trees such that every vertex has degree 1, 2 or 4).

#### 2 The Randić matrix

Let  $\mathcal{G}$  be a collection of graphs with n vertices. We would like, for a given graph  $G \in \mathcal{G}$ , to describe the set

$$[G]_{\chi} = \{ H \in \mathcal{G} : \chi(H) = \chi(G) \}$$

As we mentioned in the introduction, the idea is to associate to  $\mathcal{G}$  a  $\mathbb{Q}$ -linear map  $\rho : \mathbb{Q}^m \to \mathbb{Q}^k$  (for some integers k, m depending on  $\mathcal{G}$ ). The kernel of  $\rho$  contains the information needed to characterize  $[G]_{\chi}$  in terms of a system of linear equations. Heading in this direction we introduce some notation.

Let  $\mathbb{D}(\mathcal{G})$  be the set of vertex degrees of graphs in  $\mathcal{G}$  and

$$\mathbb{X}(\mathcal{G}) = \{i \cdot j : i, j \in \mathbb{D}(\mathcal{G}) \text{ and } i < j\}.$$

Given  $x \in \mathbb{X}(\mathcal{G})$ , let  $x = p_1^{\alpha_1} \cdots p_r^{\alpha_r}$  be the prime decomposition and define  $E(x) = \{i \in \{1, \dots, r\} : \alpha_i \text{ is even}\}$  and  $O(x) = \{i \in \{1, \dots, r\} : \alpha_i \text{ is odd}\}$ . So we can express every  $x \in \mathbb{X}(\mathcal{G})$  as  $x = \prod_{i \in E(x)} p_i^{\alpha_i} \cdot \prod_{i \in O(x)} p_i^{\alpha_i}$ . If  $y = \prod_{j \in E(y)} q_j^{\beta_j} \cdot \prod_{j \in O(y)} q_j^{\beta_j} \in \mathbb{X}(\mathcal{G})$  we define an

equivalence relation over  $\mathbb{X}(\mathcal{G})$  as follows:

$$x \sim y \Longleftrightarrow \{p_i : i \in O(x)\} = \{q_j : j \in O(y)\}$$

Denote by  $\overline{\mathbb{X}(\mathcal{G})}$  the quotient set of  $\mathbb{X}(\mathcal{G})$  modulo this equivalence relation and by [x] the equivalence class of  $x \in \mathbb{X}(\mathcal{G})$ .

Finally, we will define two functions  $\Psi : \overline{\mathbb{X}(\mathcal{G})} \longrightarrow \mathbb{R}$  and  $\Gamma : \mathbb{X}(\mathcal{G}) \longrightarrow \mathbb{N}$  such that  $\sqrt{x} = \Gamma(x)\Psi([x])$  for  $x \in \mathbb{X}(\mathcal{G})$ . Let  $\Psi$  be defined as follows:

$$\Psi\left([x]\right) = \begin{cases} \sqrt{\prod_{i \in O(x)} p_i} & \text{if } O(x) \neq \emptyset\\ 1 & \text{if } O(x) = \emptyset \end{cases}$$

and let  $\Gamma$  be defined by

$$\Gamma\left(x\right) = \prod_{i \in E(x)} p_i^{\frac{\alpha_i}{2}} \times \prod_{i \in O(x)} p_i^{\frac{\beta_i - 1}{2}}$$

For instance,  $\Gamma(2^3 \cdot 3^4 \cdot 5) = 2 \cdot 3^2$ . In other words,  $\Gamma(x)$  is the part of x that can be taken out as an integer of the square root of x.

The following lemma is probably known, but we will sketch its proof for the sake of completeness.

**Lemma 2.1.**  $\{\Psi([x])\}_{[x]\in\overline{\mathbb{X}(\mathcal{G})}}$  is a linearly independent set over  $\mathbb{Q}$ .

*Proof.* Let  $p_1, \ldots, p_r$  be the set of all different prime numbers appearing as odd powers in the prime decomposition of each  $[x] \in \overline{\mathbb{X}(\mathcal{G})}$ . Consider the following tower of field extensions  $K_0 \subset K_1 \subset \cdots \subset K_r$ 

where for each  $1 \leq i \leq r$ ,  $K_i = K_{i-1}(\sqrt{p_i})$  (the field obtained from  $K_{i-1}$  by adjoining  $\sqrt{p_i}$ ). Since  $\{1, \sqrt{p_i}\}$  is a basis of  $K_i$  over  $K_{i-1}$  then the set

$$\mathbb{B} = \{z_1 \cdot z_2 \cdot \cdots \cdot z_r : z_i \in \{1, \sqrt{p_i}\}\}$$

forms a basis of  $K_r$  over  $K_0 = \mathbb{Q}$ . But clearly  $\{\Psi([x])\}_{[x]\in\overline{\mathbb{X}(\mathcal{G})}} \subseteq \mathbb{B}$  which implies that  $\{\Psi([x])\}_{[x]\in\overline{\mathbb{X}(\mathcal{G})}}$  is a linearly independent set over  $\mathbb{Q}$ .

Now we have all we need to show a key lemma

**Lemma 2.2.** Let  $\mathcal{G}$  be a collection of graphs with n vertices  $(n \geq 5)$  and  $\chi : \mathcal{G} \longrightarrow \mathbb{R}$  the Randić function. Suppose there is  $u \in \mathbb{X}(\mathcal{G})$  such that  $O(u) = \emptyset$  (i.e. u is a perfect square) and let  $k = l.c.m. \{\mathbb{X}(\mathcal{G})\}$ . Then for every  $G \in \mathcal{G}$ ,

$$k (2\chi(G) - n) = \sum_{ij \in [u]} A_{ij} m_{ij}(G) - \sum_{\substack{[x] \in \overline{\mathbb{X}(G)} \\ [x] \neq [u]}} \sum_{ij \in [x]} B_{ij} m_{ij}(G) + \sum_{\substack{[x] \in \overline{\mathbb{X}(G)} \\ [x] \neq [u]}} \left( \sum_{ij \in [x]} C_{ij} m_{ij}(G) \right) \Psi([x])$$

where  $A_{ij} = \frac{k(2\Gamma(ij)-i-j)}{ij} \in \mathbb{Z}, B_{ij} = \frac{k(i+j)}{ij} \in \mathbb{N}$  and  $C_{ij} = \frac{2k\Gamma(ij)}{ij} \in \mathbb{N}$ .

*Proof.* For  $G \in \mathcal{G}$  we know from (2) that

$$\chi(G) = \frac{n(G)}{2} - \frac{1}{2} \sum_{1 \le i < j \le n-1} \left(\frac{1}{j} + \frac{1}{i}\right) m_{ij}(G) + \sum_{1 \le i < j \le n-1} \frac{\sqrt{ij}}{ij} m_{ij}(G)$$
(3)

and

$$\sum_{1 \le i < j \le n-1} \frac{\sqrt{ij}}{ij} m_{ij}(G) = \sum_{[x] \in \overline{\mathbb{X}(\mathcal{G})}} \left[ \sum_{ij \in [x]} \frac{\Gamma(ij)}{ij} m_{ij}(G) \right] \Psi([x])$$
$$= \sum_{ij \in [u]} \frac{\Gamma(ij)}{ij} m_{ij}(G) + \sum_{\substack{[x] \in \overline{\mathbb{X}(\mathcal{G})} \\ [x] \neq [u]}} \left[ \sum_{ij \in [x]} \frac{\Gamma(ij)}{ij} m_{ij}(G) \right] \Psi([x])$$

By substituting this last equation in (3) and then multiplying by 2k we get the result.  $\Box$ 

The integer coefficients  $A_{ij}, B_{ij}$  and  $C_{ij}$  that appeared in Lemma 2.2 gives a natural way to associate a linear map  $\rho_{\mathcal{G}} : \mathbb{Q}^{|\mathbb{X}(\mathcal{G})|} \longrightarrow \mathbb{Q}^{|\overline{\mathbb{X}(\mathcal{G})}|}$  to every collection  $\mathcal{G}$  of graphs with a fixed number of vertices as follows:

$$\rho_{\mathcal{G}} = \begin{pmatrix} \begin{bmatrix} A_{ij} \end{bmatrix}_{ij \in [u]} & \begin{bmatrix} -B_{ij} \end{bmatrix}_{ij \in [x_1]} & \begin{bmatrix} -B_{ij} \end{bmatrix}_{ij \in [x_2]} & \cdots & \begin{bmatrix} -B_{ij} \end{bmatrix}_{ij \in [x_r]} \\ \begin{bmatrix} 0 \end{bmatrix}_{ij \in [u]} & \begin{bmatrix} C_{ij} \end{bmatrix}_{ij \in [x_1]} & \begin{bmatrix} 0 \end{bmatrix}_{ij \in [x_2]} & \cdots & \begin{bmatrix} 0 \end{bmatrix}_{ij \in [x_r]} \\ \begin{bmatrix} 0 \end{bmatrix}_{ij \in [u]} & \begin{bmatrix} 0 \end{bmatrix}_{ij \in [x_1]} & \begin{bmatrix} C_{ij} \end{bmatrix}_{ij \in [x_2]} & \cdots & \begin{bmatrix} 0 \end{bmatrix}_{ij \in [x_r]} \\ \vdots & \vdots & \vdots & \vdots \\ \begin{bmatrix} 0 \end{bmatrix}_{ij \in [u]} & \begin{bmatrix} 0 \end{bmatrix}_{ij \in [x_1]} & \begin{bmatrix} 0 \end{bmatrix}_{ij \in [x_2]} & \cdots & \begin{bmatrix} C_{ij} \end{bmatrix}_{ij \in [x_r]} \end{pmatrix}$$

where  $\overline{\mathbb{X}(\mathcal{G})} = \{[u], [x_1], \dots, [x_r]\}$  is ordered by the induced order of  $\{\Psi([x])\}_{[x]\in\overline{\mathbb{X}(\mathcal{G})}}$  and the elements in each class of  $\overline{\mathbb{X}(\mathcal{G})}$  are lexicographically ordered (viewing them as a set of two integers). We call the matrix above the *Randić matrix of*  $\mathcal{G}$ . For the sake of simplicity, when there is no danger of confusion about the collection  $\mathcal{G}$ , we will write  $\rho$  instead of  $\rho_{\mathcal{G}}$ .

**Remark 2.3.** If  $X(\mathcal{G})$  contains no perfect square numbers, then  $k(2\chi - n)(G)$  can be expressed in a simpler way as

$$k(2\chi - n)(G) = -\sum_{[x]\in\overline{\mathbb{X}(\mathcal{G})}}\sum_{ij\in[x]} B_{ij}m_{ij}(G) + \sum_{[x]\in\overline{\mathbb{X}(\mathcal{G})}} \left(\sum_{ij\in[x]} C_{ij}m_{ij}(G)\right)\Psi([x])$$

In this case, the Randić matrix of  $\mathcal{G}$  is

$$\rho_{\mathcal{G}} = \begin{pmatrix}
[-B_{ij}]_{ij\in[x_1]} & [-B_{ij}]_{ij\in[x_2]} & \cdots & [-B_{ij}]_{ij\in[x_r]} \\
[C_{ij}]_{ij\in[x_1]} & [0]_{ij\in[x_2]} & \cdots & [0]_{ij\in[x_r]} \\
[0]_{ij\in[x_1]} & [C_{ij}]_{ij\in[x_2]} & \cdots & [0]_{ij\in[x_r]} \\
\vdots & \vdots & \vdots \\
[0]_{ij\in[x_1]} & [0]_{ij\in[x_2]} & \cdots & [C_{ij}]_{ij\in[x_r]}
\end{pmatrix}$$
(4)

which determines a  $\mathbb{Q}$ -linear map  $\rho_{\mathcal{G}} : \mathbb{Q}^{|\mathbb{X}(\mathcal{G})|} \longrightarrow \mathbb{Q}^{|\overline{\mathbb{X}(\mathcal{G})}|+1}$ 

It is convenient at this point to redefine the notion of the Randić structure of a graph (see the introduction) so that it includes the order of the columns of the Randić matrix.

**Definition 2.4.** The Randić structure of a graph G with n vertices, denoted by R(G), is defined as the ordered  $|\mathbb{X}(\mathcal{G})|$ -tuple

$$R(G) = (m_{ij})_{ij \in \mathbb{X}(\mathcal{G})} \in \mathbb{Q}^{|\mathbb{X}(\mathcal{G})|}$$

where  $ij \in \mathbb{X}(\mathcal{G})$  varies in the same order as the columns of  $\rho_{\mathcal{G}}$ .

Notice that R(G) is empty when all components in G are regular graphs.

In this way we have the Randić structure function  $\mathcal{G} \xrightarrow{R} \mathbb{Q}^{|\mathbb{X}(\mathcal{G})|}$ , defined by  $R(G) = (m_{ij})_{ij \in \mathbb{X}(\mathcal{G})}$ . The content of Lemma 2.2 is that, for a  $G \in \mathcal{G}$ ,  $\chi(G)$  can be expressed in terms of the Randić matrix of  $\mathcal{G}$  as follows

$$k(2\chi(G) - n) = \rho(R(G)) \cdot (\Psi([u]), \Psi([x_1]), \dots, \Psi([x_r]))$$
(5)

where  $\cdot$  means the usual inner product.

Now we are ready to present a basic representation of  $[G]_{\chi}$  for a general collection  $\mathcal{G}$  of graphs.

**Theorem 2.5.** Let  $\mathcal{G}$  be a collection of graphs with n vertices and let  $\rho_{\mathcal{G}}$  be the linear map associate with  $\mathcal{G}$ . Then the following holds for every  $G, H \in \mathcal{G}$ .

$$\chi(G) = \chi(H) \Longleftrightarrow R(G) - R(H) \in \ker \rho_{\mathcal{G}}$$

*Proof.* For every  $G \in \mathcal{G}$ , we know from equation (5) that

$$k(2\chi - n)(G) = \rho(R(G)) \cdot (\Psi[u], \Psi[x_1], \dots, \Psi[x_r])$$

From this it follows that

$$\chi(G) = \chi(H) \iff \rho(R(G) - R(H)) \cdot (\Psi[u], \Psi[x_1], \dots, \Psi[x_r]) = 0.$$

To finish the proof we recall that by Lemma 2.1 the  $\Psi[x]$ 's are all independent over  $\mathbb{Q}$ .  $\Box$ 

**Example 2.6.** Consider the collection of chemical graphs C = C(n) (i.e. graphs in which no vertex has degree greater than 4) with n vertices. Then

$$\mathbb{X}(\mathcal{C}) = \{1 \cdot 2, 1 \cdot 3, 1 \cdot 4, 2 \cdot 3, 2 \cdot 4, 3 \cdot 4\}$$

and

$$\overline{\mathbb{X}(\mathcal{C})} = \{ [1 \cdot 4], [1 \cdot 2], [1 \cdot 3], [2 \cdot 3] \}$$

Let us calculate the entries of  $\rho_{\mathcal{C}}$ 

$$A_{14} = \frac{24(2 \cdot 2 - 5)}{4} = -6 \qquad B_{12} = \frac{24(1+2)}{2} = 36 \qquad B_{24} = \frac{24(2+4)}{8} = 18$$
$$B_{13} = \frac{24(1+3)}{3} = 32 \qquad B_{34} = \frac{24(3+4)}{12} = 14 \qquad B_{23} = \frac{24(2+3)}{6} = 20$$
$$C_{12} = \frac{2 \cdot 24 \cdot 1}{2} = 24 \qquad C_{24} = \frac{2 \cdot 24 \cdot 2}{8} = 12 \qquad C_{13} = \frac{2 \cdot 24 \cdot 1}{3} = 16$$
$$C_{34} = \frac{2 \cdot 24 \cdot 2}{12} = 8 \qquad C_{23} = \frac{2 \cdot 24 \cdot 1}{6} = 8$$

and the matrix is

It can be easily checked that ker  $(\rho_{\mathcal{C}}) = \langle (-2, 0, 0, 3, -6, 0), (0, 1, -2, 0, 0, 0) \rangle$ , where  $\langle X \rangle$  denotes the subspace generated by X.

Now fix  $G \in C$  and let  $R(G) = (m_{14}(G), m_{12}(G), m_{24}(G), m_{13}(G), m_{34}(G), m_{23}(G))$ . By Theorem 2.5

$$[G]_{\chi} = \{ H \in \mathcal{C} : R(H) - R(G) \in \langle (-2, 0, 0, 3, -6, 0), (0, 1, -2, 0, 0, 0) \rangle \}$$

In other words,  $H \in [G]_{\gamma}$  if and only if there exist integers a, b such that

$$m_{14}(H) = m_{14}(G) - 2a$$
  

$$m_{12}(H) = m_{12}(G) + b$$
  

$$m_{24}(H) = m_{24}(G) - 2b$$
  

$$m_{13}(H) = m_{13}(G) + 3a$$
  

$$m_{34}(H) = m_{34}(G) - 6a$$
  

$$m_{23}(H) = m_{23}(G)$$

We end this section with a couple of remarks in order to clarify the dependency of the Randić matrix on the collection  $\mathcal{G}$  and also the role played by the identity (2) in our definition of  $\rho$ .

**Remark 2.7.** (i) The order we have been using for the columns of a Randić matrix and for the Randić structure of a graphs does not really depend on the collection  $\mathcal{G}$ . In fact, it only depends on n.

(ii) Given two collections  $\mathcal{G}' \subset \mathcal{G}$ , it is not difficult to show that  $\rho_{\mathcal{G}'}$  can be easily computed from  $\rho_{\mathcal{G}}$ . In fact  $\rho_{\mathcal{G}'}$  is obtained by deleting the columns of  $\rho_{\mathcal{G}}$  which do not correspond to elements of  $\mathbb{X}(\mathcal{G}')$  and then multiplying by an appropriate integer (determined by the m.c.m used to define  $\rho_{\mathcal{G}}$  and  $\rho_{\mathcal{G}'}$ ).

(iii) There are other identities like (2) that could be used to associate a matrix to  $\chi$ . The advantage of using (2) is that the number of variables used is minimal.

### 3 The Randić structure property

As we said in the introduction, perhaps the simplest situation, regarding the problem of describing  $[G]_{\chi}$ , is when  $[G]_{\chi}$  consist merely of those  $H \in \mathcal{G}$  such that R(H) = R(G). In this section we will present examples of such collections. We recall a concept already mentioned in the introduction.

**Definition 3.1.** A collection  $\mathcal{G}$  of graphs with a fixed number of vertices has the Randić structure property (RSP) if for all  $G, G' \in \mathcal{G}$ 

$$\chi(G) = \chi(G') \iff R(G) = R(G')$$

Notice that the implication from right to left always holds. In other words, if  $\mathcal{G}$  has RSP then for every  $G \in \mathcal{G}$ 

$$[G]_{\gamma} = \{ H \in \mathcal{G} : m_{ij}(H) = m_{ij}(G) \text{ for all } i < j \in \mathbb{D}(\mathcal{G}) \}$$

Let us define  $\Delta_{\mathcal{G}} : \mathcal{G} \times \mathcal{G} \longrightarrow \mathbb{Q}^{|\mathbb{X}(\mathcal{G})|}$  as  $\Delta(G, H) = R(G) - R(H)$  for every  $G, H \in \mathcal{G}$ . From Theorem 2.5 we immediately get the following

**Proposition 3.2.** Let  $\mathcal{G}$  be a collection of graphs with n vertices and  $\rho_{\mathcal{G}}$  its associated linear map. The following conditions are equivalent:

- 1.  $\mathcal{G}$  has the Randić structure property;
- 2.  $Im(\Delta_{\mathcal{G}}) \cap \ker \rho_{\mathcal{G}} = (0).$

In particular, if  $\rho_{\mathcal{G}}$  is one to one then  $\mathcal{G}$  has the RSP. Our next result characterizes collections  $\mathcal{G}$  such that  $\rho_{\mathcal{G}}$  is one to one.

**Proposition 3.3.** Let  $\mathcal{G}$  be a collection of graphs with n vertices and  $\rho_{\mathcal{G}}$  its associated linear map. The following conditions are equivalent:

- 1.  $\rho_{\mathcal{G}}$  is one to one;
- 2.  $|\mathbb{X}(\mathcal{G})| = \left|\overline{\mathbb{X}(\mathcal{G})}\right|$ .

*Proof.* If there exists a perfect square number  $u \in \mathbb{X}(\mathcal{G})$  then dim  $(Im\rho_{\mathcal{G}}) = \left|\overline{\mathbb{X}(\mathcal{G})}\right|$  since  $\rho_{\mathcal{G}}$  has exactly  $\left|\overline{\mathbb{X}(\mathcal{G})}\right|$  independent rows. Consequently, dim  $(\ker \rho_{\mathcal{G}}) = |\mathbb{X}(\mathcal{G})| - \left|\overline{\mathbb{X}(\mathcal{G})}\right|$  and the result follows.

Let us assume then that  $\mathbb{X}(\mathcal{G})$  has no perfect squares and so its Randić matrix has the form (4). Note that  $\left|\overline{\mathbb{X}(\mathcal{G})}\right| \leq \dim (Im\rho_{\mathcal{G}}) \leq \left|\overline{\mathbb{X}(\mathcal{G})}\right| + 1$  which implies

$$|\mathbb{X}(\mathcal{G})| - \left|\overline{\mathbb{X}(\mathcal{G})}\right| - 1 \le \dim\left(\ker\rho_{\mathcal{G}}\right) \le |\mathbb{X}(\mathcal{G})| - \left|\overline{\mathbb{X}(\mathcal{G})}\right|$$
(6)

 $1. \Rightarrow 2. \text{ Suppose that } |\mathbb{X}(\mathcal{G})| > \left|\overline{\mathbb{X}(\mathcal{G})}\right|. \text{ If } |\mathbb{X}(\mathcal{G})| - \left|\overline{\mathbb{X}(\mathcal{G})}\right| \ge 2 \text{ then, by (6), dim (ker } \rho_{\mathcal{G}}) \ge 1 \text{ and so } \ker \rho_{\mathcal{G}} \neq (0).$ 

Next we show that if  $\mathbb{X}(\mathcal{G})$  has no perfect squares then  $|\mathbb{X}(\mathcal{G})| - |\overline{\mathbb{X}(\mathcal{G})}| \neq 1$ . If  $|\mathbb{X}(\mathcal{G})| - |\overline{\mathbb{X}(\mathcal{G})}| = 1$  then there exists  $i \cdot j$ ,  $r \cdot s \in \mathbb{X}(\mathcal{G})$  such that  $\{i, j\} \neq \{r, s\}$  and  $i \cdot j \sim r \cdot s$ . Consequently, we have expressions of the form

$$i \cdot j = \prod_{k=1}^{u} p_k^{e_k} \times \prod_{k=1}^{v} q_k^{o_k} \qquad r \cdot s = \prod_{k=1}^{u} p_k^{e'_k} \times \prod_{k=1}^{v} q_k^{o'_k}$$

where for all k,  $e_k$  and  $e'_k$  are even (possibly zero) natural numbers and  $o_k$  and  $o'_k$  are odd natural numbers. Let

$$i = \prod_{k=1}^{u} p_k^{\alpha_k} \times \prod_{k=1}^{v} q_k^{\beta_k} \qquad j = \prod_{k=1}^{u} p_k^{\gamma_k} \times \prod_{k=1}^{v} q_k^{\delta_k}$$
$$r = \prod_{k=1}^{u} p_k^{\alpha'_k} \times \prod_{k=1}^{v} q_k^{\beta'_k} \qquad s = \prod_{k=1}^{u} p_k^{\gamma'_k} \times \prod_{k=1}^{v} q_k^{\delta'_k}$$

where for each k,

$$\alpha_k + \gamma_k = e_k \qquad \beta_k + \delta_k = o_k$$

$$\alpha'_k + \gamma'_k = e'_k \qquad \beta'_k + \delta'_k = o'_k$$
(7)

We claim that  $i \cdot r \sim j \cdot s$ . In fact,

$$i \cdot r = \prod_{k=1}^u p_k^{\alpha_k + \alpha'_k} \times \prod_{k=1}^v q_k^{\beta_k + \beta'_k} \qquad j \cdot s = \prod_{k=1}^u p_k^{\gamma_k + \gamma'_k} \times \prod_{k=1}^v q_k^{\delta_k + \delta'_k}$$

where by (7)

$$(\alpha_k + \alpha'_k) + (\gamma_k + \gamma'_k) = e_k + e'_k$$
  
$$(\beta_k + \beta'_k) + (\delta_k + \delta'_k) = o_k + o'_k$$

Since  $e_k + e'_k$  and  $o_k + o'_k$  are even, we deduce that

$$(\alpha_k + \alpha'_k)$$
 is even  $\Leftrightarrow$   $(\gamma_k + \gamma'_k)$  is even  
 $(\beta_k + \beta'_k)$  is even  $\Leftrightarrow$   $(\delta_k + \delta'_k)$  is even

Consequently,  $i \cdot r \sim j \cdot s$ .

Now, since  $|\mathbb{X}(\mathcal{G})| - \left|\overline{\mathbb{X}(\mathcal{G})}\right| = 1$  we must have r = j or i = s. Assume r = j. Then for all  $k, \alpha'_k = \gamma_k$  and  $\beta'_k = \delta_k$ , which implies by (7)

 $\alpha_k + \alpha'_k = e_k \qquad \beta_k + \beta'_k = o_k$  $\alpha'_k + \gamma'_k = e'_k \qquad \beta'_k + \delta'_k = o'_k$ 

Hence,

$$\alpha_k$$
 is even  $\Leftrightarrow \gamma'_k$  is even  
 $\beta_k$  is even  $\Leftrightarrow \delta'_k$  is even

This clearly implies that  $\alpha_k + \gamma'_k$  and  $\beta_k + \delta'_k$  are even for all k. Finally, since

$$i\cdot s = \prod_{k=1}^u p_k^{\alpha_k+\gamma'_k} \times \prod_{k=1}^v q_k^{\beta_k+\delta'_k}$$

we conclude that  $i \cdot s$  is a perfect square, but this is a contradiction. Similarly, i = s implies  $r \cdot j$  is a perfect square which also yields a contradiction.

 $2. \Rightarrow 1$ . This is an immediate consequence of (6).

An application of Propositions (3.2) and (3.3) gives the following examples

**Example 3.4.** Let  $\mathcal{P}_n$  be the collection of all graphs with n vertices such that all vertex degrees are prime numbers or 1. Then clearly the condition  $|\mathbb{X}(\mathcal{P}_n)| = \left|\overline{\mathbb{X}(\mathcal{P}_n)}\right|$  is satisfied for all n. Hence  $\mathcal{P}_n$  has the Randić structure property. In particular, the collection  $\mathcal{B}_n$  of benzenoid systems with n vertices has RSP since it is a subset of  $\mathcal{P}_n$ .

**Example 3.5.** Let  $\mathcal{G} = \mathcal{R}_{\delta}(n)$  be the set of all branch regular trees of degree  $\delta$  and n vertices (see [13]). We recall that  $G \in \mathcal{R}_{\delta}(n)$  if and only if all branching vertices of G have degree  $\delta$ . It can be easily checked that for all values of  $\delta$ , except when  $\delta$  is a perfect square or an odd power of 2,  $|\mathbb{X}(\mathcal{R}_{\delta}(n))| = |\overline{\mathbb{X}(\mathcal{R}_{\delta}(n))}|$ . It follows that for these values of  $\delta$ ,  $\mathcal{R}_{\delta}(n)$  has RSP. In section §5 we will present examples showing that  $\mathcal{R}_{\delta}(n)$  has not the RSP for  $\delta$  a perfect square or an odd power of 2.

Up to now all examples of RSP collections  $\mathcal{G}$  have ker  $(\rho_{\mathcal{G}}) = (0)$ . As we will see in our next example this is not always the case.

**Example 3.6.** Let  $S_4 = S_4(n)$  be the collection of all starlike trees of degree 4 and n vertices. Recall that  $T \in S_4$  if and only if T has a unique branching vertex of degree 4. We have  $\mathbb{X}(S_4) = \{1 \cdot 2, 1 \cdot 4, 2 \cdot 4\}$ . Since  $1 \cdot 2$  and  $2 \cdot 4$  are equivalent, then  $\overline{\mathbb{X}(S_4)} = \{[1 \cdot 4], [1 \cdot 2]\}$  and k = 8. The associated matrix  $\rho = \rho_{S_4}$  is

$$\rho = \left(\begin{array}{ccc}
-2 & -12 & -6\\
0 & 8 & 4
\end{array}\right)$$
(8)

where the columns correspond to the pair  $1 \cdot 4$ ,  $1 \cdot 2$  and  $2 \cdot 4$  in that order. The kernel of  $\rho$  can be easily shown to be generated by (0, -2, 1). Now, for every  $S \in S_4$  the following relations hold

$$m_{12}(S) = m_{24}(S) = 4 - m_{14}(S)$$

Consequently,

$$\Delta_{\mathcal{S}_4}(S,S') = R(S) - R(S') = (m_{14} - m'_{14}, m_{12} - m'_{12}, m_{24} - m'_{24}) \subseteq \langle (1, -1, -1) \rangle$$

Since  $\langle (1, -1, -1) \rangle \cap \ker(\rho_{S_4}) = (0)$ , we deduce that  $Im(\Delta_{S_4}) \cap \ker(\rho_{S_4}) = (0)$ . It follows by Proposition 3.2, that  $S_4$  has RSP.

#### 4 Decomposition of a collection of graphs into disjoint

#### **RSP** subcollections

Even though collections with RSP might seem hard to find, we will show in this section a fairly general method to generate them. Furthermore, we will show that every collection  $\mathcal{G}$  can be decomposed as a disjoint union of RSP subcollections.

Let  $\mathcal{G}$  be a collection of graphs with a fixed number of vertices and  $\rho_{\mathcal{G}} : \mathbb{Q}^{|X(\mathcal{G})|} \longrightarrow \mathbb{Q}^{|\overline{X(\mathcal{G})}|}$ its associated  $\mathbb{Q}$ -linear map. It is clear from Proposition 3.2 that a subcollection  $\mathcal{G}' \subseteq \mathcal{G}$  has the RSP if and only if  $\rho_{\mathcal{G}}$  is one-to-one in  $\{R(G) : G \in \mathcal{G}'\}$ . The basic idea for getting the above mentioned decomposition of  $\mathcal{G}$  is first to decompose  $\mathbb{Q}^{|X(\mathcal{G})|}$  into disjoint pieces where  $\rho_{\mathcal{G}}$  is one-to-one and then pull back with  $R^{-1}$  such decomposition into  $\mathcal{G}$ .

**Lemma 4.1.** Let  $\rho : \mathbb{Q}^m \longrightarrow \mathbb{Q}^k$  and  $\sigma : \mathbb{Q}^m \longrightarrow \mathbb{Q}^{m-k}$  be  $\mathbb{Q}$ -linear maps and consider the  $\mathbb{Q}$ -linear map  $(\rho, \sigma) : \mathbb{Q}^m \longrightarrow \mathbb{Q}^m$ , defined by  $(\rho, \sigma)(Z) = (\rho(Z), \sigma(Z))$  for every  $Z \in \mathbb{Q}^m$ . If  $(\rho, \sigma)$  is invertible then ker  $(\sigma)$  is a complementary direct summand of ker $(\rho)$  in  $\mathbb{Q}^m$ .

Proof. It is clear that  $(0) = \ker((\rho, \sigma)) = \ker(\rho) \cap \ker(\sigma)$ . Now choose  $Z \in \mathbb{Q}^m$ . Since  $(\rho, \sigma)$  is onto there exists  $Y \in \mathbb{Q}^m$  such that  $(\rho, \sigma)(Y) = (0, \sigma(Z))$ . Consequently,  $Y \in \ker(\rho)$ ,  $Z - Y \in \ker(\sigma)$  and Z = Y + (Z - Y).

**Proposition 4.2.** Let  $\sigma : \mathbb{Q}^{|X(\mathcal{G})|} \longrightarrow \mathbb{Q}^{|X(\mathcal{G})| - |\overline{X(\mathcal{G})}|}$  be a  $\mathbb{Q}$ -linear map such that  $(\rho_{\mathcal{G}}, \sigma)$  is invertible and  $\mathcal{G}_0$  a subcollection of  $\mathcal{G}$ . If  $\sigma \circ \Delta_{\mathcal{G}_0} = (0)$  then  $\mathcal{G}_0$  has the RSP.

*Proof.* Let  $G, G' \in \mathcal{G}_0$  and suppose that  $\chi(G) = \chi(G')$ . Then by Theorem 2.5,  $R(G) - R(G') \in \ker(\rho_{\mathcal{G}})$ . On the other hand  $R(G) - R(G') \in \ker(\sigma)$  since  $\sigma \circ \Delta_{\mathcal{G}_0} = (0)$ . It follows from Lemma 4.1 that R(G) = R(G') and so  $\mathcal{G}_0$  has the RSP.  $\Box$ 

**Theorem 4.3.** Let  $\mathcal{G}$  be a collection of graphs with a fixed number of vertices. Then  $\mathcal{G}$  can be decomposed into pairwise disjoint RSP subcollections.

Proof. Choose a complementary direct summand W of ker  $(\rho_{\mathcal{G}})$  in  $\mathbb{Q}^{|X(\mathcal{G})|}$ . Consider the projection  $\pi : \mathbb{Q}^{|X(\mathcal{G})|} = W \oplus \ker(\rho_{\mathcal{G}}) \longrightarrow \ker(\rho_{\mathcal{G}}) \cong \mathbb{Q}^{|X(\mathcal{G})|-|\overline{X(\mathcal{G})}|}$ . Clearly  $(\rho_{\mathcal{G}}, \pi) : \mathbb{Q}^{|X(\mathcal{G})|} \longrightarrow \mathbb{Q}^{|X(\mathcal{G})|}$  is an invertible  $\mathbb{Q}$ -linear map. For each  $k \in Im(\pi \circ R)$  define  $\mathcal{G}_k = \{G \in \mathcal{G} : R(G) \in W + k\}$ . Then for every  $G, G' \in \mathcal{G}_k$  we have  $R(G) - R(G') \in W$  and consequently,  $\pi(R(G) - R(G')) = 0$ . Hence  $\pi \circ \Delta_{\mathcal{G}_k} = (0)$  and so, by Proposition 4.2,  $\mathcal{G}_k$  has the RSP. Finally, it is clear that the  $\mathcal{G}_k$  's are pairwise disjoint and  $\mathcal{G} = \bigcup_{k \in Im(\pi \circ R)} \mathcal{G}_k$ .  $\Box$ 

Let us look at a concrete example. Consider the collection  $\mathcal{R}_{\delta}(n)$  of branch regular graphs of degree  $\delta$  and n vertices. We analyze the case where  $\delta = q^2$  is a perfect square. Then  $\mathbb{X}(\mathcal{R}_{\delta}) = \{1 \cdot 2, 1 \cdot \delta, 2 \cdot \delta\}, \overline{\mathbb{X}(\mathcal{R}_{\delta})} = \{[1 \cdot \delta], [1 \cdot 2]\}$  and  $k = 2\delta$ . The associated matrix is

$$\rho = \begin{pmatrix} -2(q-1)^2 & -3q^2 & -(q^2+2) \\ 0 & 2q^2 & 2q \end{pmatrix}$$

Recall that the columns correspond to the following pairs  $1 \cdot \delta, 1 \cdot 2, 2 \cdot \delta$  in this order. Consider the following completion of  $\rho$ 

$$(\rho, \sigma) = \begin{pmatrix} -2(q-1)^2 & -3q^2 & -(q^2+2) \\ 0 & 2q^2 & 2q \\ 1 & 1 & 0 \end{pmatrix}$$

where  $\sigma : \mathbb{Q}^3 \longrightarrow \mathbb{Q}$  is defined by  $\sigma(x_{1\delta}, x_{12}, x_{2\delta}) = x_{1\delta} + x_{12}$ . Notice that  $(\rho, \sigma)$  is invertible. For a given a graph G, we denote by  $k_i = k_i(G)$  the number of vertices of G of degree i. For each positive integer  $r \ge 1$  consider the subcollection of  $\mathcal{R}_{\delta}$ 

$$\mathcal{R}^{r}_{\delta} = \{ G \in \mathcal{R}_{\delta} : k_{\delta}(G) = r \}$$
(9)

Since for every  $G \in \mathcal{R}_{\delta}$ ,  $m_{12}(G) + m_{1\delta}(G) = k_1(G) = (\delta - 2) k_{\delta}(G) + 2$ , we deduce that  $m_{12}(G) + m_{1\delta}(G) = (\delta - 2) r + 2$  for every  $G \in \mathcal{R}_{\delta}^r$ . Consequently, if  $G, G' \in \mathcal{R}_{\delta}^r$ 

$$\sigma \left( R \left( G \right) - R \left( G' \right) \right) = \sigma \left( \left( m_{1\delta} - m'_{1\delta}, m_{12} - m'_{12}, m_{2\delta} - m'_{2\delta} \right) \right) \\ = \left( m_{1\delta} - m'_{1\delta} \right) + \left( m_{12} - m'_{12} \right) = 0$$

In other words,  $\sigma \circ \Delta_{\mathcal{R}^r_{\delta}} = (0)$ . It follows from Proposition 4.2 that  $\mathcal{R}^r_{\delta}$  has the RSP for every r. Moreover, an easy induction shows that  $k_{\delta}$ .  $(\delta - 1) \leq n - 2$ . Therefore

$$\mathcal{R}_{\delta} = \bigcup_{r=1}^{\left[\frac{n-2}{\delta-1}\right]} \mathcal{R}_{\delta}^{r} \tag{10}$$

is a disjoint union of RSP subcollections of  $\mathcal{R}_{\delta}$ .

Note that  $\mathcal{R}^1_{\delta}$  is the set of all starlike trees of degree  $\delta$ . In particular, Example 3.6 can be deduced from here since  $\mathcal{S}_4 = \mathcal{R}^1_4$ .

The case in which  $\delta$  is an odd power of 2 is similar.

We will present one more example. It will show that such decompositions into RSP pieces can be more complex than in the previous example. In particular, they could depend on more than one parameter (due to the dimension of the kernel of the Randić matrix).

Consider the collection of all chemical graphs C with n vertices. We have already computed in example 2.6 its Randić matrix  $\rho$ . Consider the following completion of  $\rho$ 

$$(\rho, \sigma) = \begin{pmatrix} -6 & -36 & -18 & -32 & -14 & -20 \\ 0 & 24 & 12 & 0 & 0 & 0 \\ 0 & 0 & 0 & 16 & 8 & 0 \\ 0 & 0 & 0 & 0 & 0 & 8 \\ 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 \end{pmatrix}$$

where  $\sigma : \mathbb{Q}^6 \longrightarrow \mathbb{Q}^2$  is defined by  $\sigma(x_{14}, x_{12}, x_{24}, x_{13}, x_{34}, x_{23}) = (x_{14} + x_{12} + x_{13}, x_{24})$ . For each pair of integers  $0 \le r \le n$  and  $0 \le s \le n$ , let

$$C_{r,s} = \{G \in C : m_{12}(G) + m_{13}(G) + m_{14}(G) = r \& m_{24}(G) = s\}$$

Proposition 4.2 says that each  $C_{r,s}$  has the RSP (notice that some of the  $C_{r,s}$ 's are empty). Finally, it is clear that these subcollections form a partition of C.

## 5 A finer analysis of $\mathcal{R}_4$

In this section we will see how the ideas presented in previous sections can be used to make a finer analysis of the collection of branch regular trees. We will restrict our analysis to  $\mathcal{R}_4$ . But before doing that, we will present some examples showing that  $\mathcal{R}_4$  and  $\mathcal{R}_8$  does not have the RSP. The pairs of trees shown in figure 1 and 2 are easily seen to have equal  $\chi$  and different Randić structures.

#### figure 1

#### figure 2

The key fact for all our analysis is the following lemma. It tells, in terms of the number of branching vertices, where we have to look to find two trees in  $\mathcal{R}_4$  with equal  $\chi$ . Notice the crucial role played by the equations of the kernel of  $\rho_{\mathcal{R}_4}$ .

**Lemma 5.1.** Let  $T, T' \in \mathcal{R}_4$  with  $\chi(T) = \chi(T')$ . Then

(a) 
$$\frac{3k_4 + 1}{4} \le k_4' \le \frac{4k_4 - 1}{3}$$

(b)

$$m_{44} - m'_{44} = 4(k_4 - k'_4)$$

where  $k_4 = k_4(T)$  denotes the number of vertices of T of degree 4 (all variable with ' correspond to T' and the others to T).

*Proof.* Let  $\mathcal{R}_4 = \mathcal{R}_4(n)$ . Since  $\mathbb{X}(\mathcal{R}_4) = \mathbb{X}(\mathcal{S}_4)$ , then we have already computed the Randić matrix  $\rho = \rho_{\mathcal{R}_4}$  in example 3.6.

$$\rho = \left(\begin{array}{ccc}
-2 & -12 & -6\\
0 & 8 & 4
\end{array}\right)$$
(11)

where the columns correspond to the pair  $1 \cdot 4$ ,  $1 \cdot 2$  and  $2 \cdot 4$  in that order. The kernel of  $\rho$  is generated by (0, -2, 1). Suppose  $\chi(T) = \chi(T')$  then by theorem 2.5 we know that

$$m_{14} = m'_{14}$$
  

$$2m_{12} + m_{24} = 2m'_{12} + m'_{24}$$
(12)

Now we start the proof of (a). For every branch regular tree we have  $m_{12} + m_{1\delta} = (\delta - 2)k_{\delta} + 2$ . Since  $m_{14} = m'_{14}$ , then

$$m_{12}' = 2k_4' + 2 - m_{14} \tag{13}$$

Let  $r(T) = 2m_{12} + m_{24}$ . By substituting (13) in (12) we get

$$m_{24}' = r(T) - 4k_4' - 4 + 2m_{14} \tag{14}$$

For every tree in  $\mathcal{R}_4$  we have that  $m'_{12} \leq m'_{24}$ . This inequality together with (12) gives the following

$$3m'_{24} \ge r(T)$$

From this and (14) we get

$$k_4' \le \frac{r(T)}{6} + \frac{m_{14}}{2} - 1$$

Since  $m_{12} + m_{14} = 2k_4 + 2$ , we immediately get

$$k_4' \le \frac{2}{3}k_4 + \frac{m_{14} + m_{24}}{6} - \frac{1}{3} \tag{15}$$

Since  $m_{14} + m_{24} + 2m_{44} = 4k_4$ , then  $m_{14} + m_{24} \le 4k_4$ . This last inequality together with (15) gives the right hand side of part (a) of our claim. The other inequality follows by symmetry.

(b) Since  $m_{14} + m_{24} + 2m_{44} = 4k_4$  and  $m_{14} = m'_{14}$ , then we have

$$4(k_4 - k'_4) = 2(m_{44} - m'_{44}) + m_{24} - m'_{24}$$
(16)

From (12) we get

$$2(k_4 - k_4') = m_{44} - m_{44}' + m_{12}' - m_{12}$$
<sup>(17)</sup>

Since  $m_{12} + m_{14} = 2k_4 + 2$  and  $m_{14} = m'_{14}$ , then we immediately get that  $m'_{12} - m_{12} = -2(k_4 - k'_4)$ . From this and (17) we are done.

Recall from §4 the partition of  $\mathcal{R}_4$  given by the  $\mathcal{R}_4^r$ 's.

**Proposition 5.2.** The following collection has the RSP for every  $r \ge 4$ 

 $\mathcal{R}_4^1 \cup \mathcal{R}_4^2 \cup \mathcal{R}_4^3 \cup \mathcal{R}_4^r$ 

*Proof.* It is a straightforward application of lemma 5.1(a). We will show the claim in three steps. First we show that  $\mathcal{R}_4^1 \cup \mathcal{R}_4^r$  has the RSP for every  $r \ge 2$ . Second we show that  $\mathcal{R}_4^2 \cup \mathcal{R}_4^r$  has the RSP for every  $r \ge 3$  and then we show that  $\mathcal{R}_4^3 \cup \mathcal{R}_4^r$  has the RSP for every  $r \ge 4$ .

(i) Let  $T \in \mathcal{R}_4^1$  and  $T' \in \mathcal{R}_4^r$  with  $\chi(T) = \chi(T')$ . Then  $k_4 = 1$  and  $k'_4 = r$ . By Lemma 5.1 we know that  $r \leq \frac{4-1}{3} = 1$ . Therefore  $T' \in \mathcal{R}_4^1$  and we are done as  $\mathcal{R}_4^1$  has the RSP. (ii) Let  $T \in \mathcal{R}_4^2$  and  $T' \in \mathcal{R}_4^r$  with  $\chi(T) = \chi(T')$ . Then from Lemma 5.1 we know that

(ii) Let  $T \in \mathcal{R}_4^2$  and  $T' \in \mathcal{R}_4^r$  with  $\chi(T) = \chi(T')$ . Then from Lemma 5.1 we know that  $r \leq \frac{8-1}{3} < 3$ . Therefore  $T' \in \mathcal{R}_4^1 \cup \mathcal{R}_4^2$  and we are done as we just saw that this collection has the RSP.

Notice that from (i) and (ii) we can conclude that  $\mathcal{R}_4^1 \cup \mathcal{R}_4^2 \cup \mathcal{R}_4^3$  has the RSP.

(iii) Let  $T \in \mathcal{R}_4^3$  and  $T' \in \mathcal{R}_4^r$  with  $\chi(T) = \chi(T')$ . Then from Lemma 5.1 we know that  $r \leq \frac{12-1}{3} < 4$ . Therefore  $T' \in \mathcal{R}_4^1 \cup \mathcal{R}_4^2 \cup \mathcal{R}_4^3$  and we are done as we just saw that this collection has the RSP.

We have already seen that the previous result is best possible, since the pair of counterexamples given in figure 2 have  $k_4$  equal to 4 and 5 respectively. But we can get nevertheless sharper results if we take into account the number of vertices. Recall that for branch regular trees we have that  $k_{\delta} \leq \frac{n-2}{\delta-1}$ . In particular, for  $\delta = 4$ , we have that  $k_4 \leq \frac{n-2}{3}$ . It is not difficult to see from 5.1 that the smallest pair of trees in  $\mathcal{R}_4$  with equal  $\chi$  and different Rhave size 19. Thus we have the following

**Proposition 5.3.** (i) For n < 19,  $\mathcal{R}_4$  has the RSP.

- (ii) For  $19 \le n < 20$ ,  $\mathcal{R}_4$  does not have the RSP but it is the union of two subcollections with the RSP, namely  $\mathcal{R}_4^1 \cup \mathcal{R}_4^2 \cup \mathcal{R}_4^3 \cup \mathcal{R}_4^4$  and  $\mathcal{R}_4^5$ .
- (iii) For  $20 \leq n < 23$ ,  $\mathcal{R}_4$  does not have the RSP but it is the union of two subcollections with the RSP, namely  $\mathcal{R}_4^1 \cup \mathcal{R}_4^2 \cup \mathcal{R}_4^3 \cup \mathcal{R}_4^4 \cup \mathcal{R}_4^6$  and  $\mathcal{R}_4^5$ .

**Remark:** Let us define an equivalence relation  $=_R$  by letting  $G =_R G'$  when R(G) = R(G'). A way of understanding the previous result is by noticing that the number of RSP pieces gives a bound to the number of  $=_R$  equivalence classes that form  $[G]_{\chi}$ . In particular, for  $G \in \mathcal{R}_4$  (with n < 23), there is at most one more Randić structure different than R(G)corresponding to a tree  $H \in \mathcal{R}_4$  with  $\chi(G) = \chi(H)$ .

Of course we can continue the previous analysis to get a quite sharp picture of  $\mathcal{R}_4$  in terms of the smallest number of RSP pieces which are necessary to cover it all. This analysis would give further information about the behavior of  $\chi$  over  $\mathcal{R}_4$ . For instance, given k, it is now easy to construct (for large enough n) a set of k trees in  $\mathcal{R}_4$  with equal  $\chi$  and different Randić structures.

## References

- O. Araujo and J. A. de la Peña, The connectivity index of a wieghted graphs, Linear Algebra Appl. 283 (1998) 171-177.
- [2] O. Araujo and J. A. de la Peña, Some bounds for the connectivity index of a chemical graph, J. Chem. Inf. Compt. Sci 38 (1998) 827-831.
- [3] O. Araujo and J. Rada, Randić index and lexicographic order, J. Math. Chem. 27 (2000) 201-212.
- [4] O. Araujo and J. Rada, Vertex and edge type relations of Randić index for chemical tress, to appear in Ars Combinatorica.
- [5] B. Bollobás and P. Erdös, Graphs of extremal weights, Ars Combinatorica 50 (1998) 225-233.
- [6] L. H. Clark and J. W. Moon, On the general Randić index for certain families of tress, Ars Combinatorica 54 (2000) 223-235.

- [7] I. Gutman, O. Miljkovic, G. Caporossi and P. Hansen, Alkanes with small and large Randić connectivity index, Chem. Phys. Lett. 306 (1999) 366-372.
- [8] I. Gutman, O. Araujo and J. Rada, An identity for Randić's connectivity index and its applications, ACH Models Chem. 137 (2000) 653-658.
- [9] I. Gutman, O. Araujo and D. Morales, Bounds for the Randić's connectivity index, to appear in J. Chem. Inf. Comput. Sci.
- [10] L.B. Kier and L.H. Hall, Molecular connectivity in chemistry and drug research, Academic Press, New York, 1976.
- [11] L.B. Kier and L.H. Hall, Molecular connectivity in structure-activity analysis, Wiley, New York, 1986.
- [12] J. Rada and O. Araujo, Higher order connectivity index of starlike tress, to appear in Discrete App. Math.
- [13] J. Rada and O. Araujo, Variation of the Randić index due to some tree operations, submitted.
- [14] M. Randić, On characterization of molecular branching, J. Am. Chem. Soc. 97 (1975) 6609-6615.



Т



T'





Т



T'







T'