

# The covering property for $\sigma$ -ideals of compact sets

Carlos E. Uzcategui A. \*

March, 1991

## Abstract

The covering property for  $\sigma$ -ideals of compact sets is an abstract version of the classical perfect set theorem for analytic sets. We will study its consequences using as a paradigm the  $\sigma$ -ideal of countable closed subsets of  $2^\omega$ .

## 1 Introduction

The study of  $\sigma$ -ideals of compact sets has been motivated by problems in analysis, and quite recently it has received a lot of attention because its connections with harmonic analysis (see [7]). The Descriptive set theoretic approach was initiated by Kechris, Louveau and Woodin in “The structure of  $\sigma$ -ideals of compact sets” ([8]) and also in “The Descriptive Set Theory of  $\sigma$ -ideals of compact sets” ([4]).

Throughout this article  $X$  will be a compact, separable metric space. By  $\mathcal{K}(X)$  we denote the collection of closed subsets of  $X$ . A subset  $I \subseteq \mathcal{K}(X)$  is called **hereditary** if

$$\text{if } K, L \in \mathcal{K}(X) ; K \in I, L \subseteq K, \text{ then } L \in I.$$

$I$  is called an **ideal**, if moreover

$$\text{if } K, L \in I, \text{ then } K \cup L \in I.$$

and  $I$  is called a  **$\sigma$ -ideal**, if in addition we have that

$$\text{if } K, K_1, K_2, \dots \in \mathcal{K}(X) \text{ are such that for all } i \ K_i \in I \text{ and } K = \bigcup K_i, \text{ then } K \in I.$$

Let us give some examples:

- (1) For each  $A \subseteq X$ , let  $\mathcal{K}(A) = \{K \in \mathcal{K}(X) : K \subseteq A\}$ .
- (2)  $\mathcal{K}_\omega(X) = \{K \in \mathcal{K}(X) : K \text{ is countable}\}$ .
- (3)  $I_{meager} = \{K \in \mathcal{K}(X) : K \text{ is meager}\}$ .
- (4) Given a Borel measure  $\mu$  over  $X$ , let

$$I_\mu = \{K \in \mathcal{K}(X) : \mu(K) = 0\}$$

- (5) Let  $R$  = Rajchman probability measure on the unit circle, i.e. those measures for which  $\hat{\mu}(n) \rightarrow 0$ , as  $|n| \rightarrow \infty$ . Let

---

\*This article is based on chapter 2 and 3 of my Caltech's Ph.D thesis under the supervision of Dr. Alexander Kechris to whom I am very grateful for his guidance and patience.

$$U_0 = \{K \in \mathcal{K}(X) : \mu(K) = 0 \text{ for all } \mu \in R\}.$$

$U_0$  are the closed sets of extended uniqueness.

(6) Let  $X = 2^\omega$ , and  $I_c =$  The  $\sigma$ -ideal of closed subsets of  $2^\omega$  that avoid a cone of Turing degrees.

Given a  $\sigma$ -ideal  $I$  of closed subsets of  $X$ , the most natural way to extend  $I$  to a  $\sigma$ -ideal of arbitrary subsets of  $X$  is as follows: Let

$$I^{ext} = \{A \subseteq X : \exists (K_n)_{n \in \omega} \text{ in } I, A \subseteq \bigcup_n K_n\}$$

$I^{ext}$  is the smallest  $\sigma$ -ideal of subsets of  $X$  extending  $I$ . A typical example is when  $I = I_{meager}$ ; the exterior extension of  $I$  is the  $\sigma$ -ideal of meager sets. Analogously the exterior extension of  $K_\omega(X)$  is the  $\sigma$ -ideal of countable sets.

In some cases, however, the exterior extension is not the natural one. For example: if  $\lambda$  is the product measure on  $2^\omega$  and  $I = I_\lambda$  then  $I^{ext}$  is not the  $\sigma$ -ideal of  $\lambda$ -measure zero sets. But this example suggests other way of extending  $I$ : Let

$$I^{int} = \{A \subseteq X : \mathcal{K}(A) \subseteq I\}$$

Clearly  $I^{int}$  is hereditary,  $I^{ext} \subseteq I^{int}$  and  $I^{int} \cap \mathcal{K}(X) = I$ . But in general  $I^{int}$  is not even an ideal.

We say that a  $\sigma$ -ideal  $I$  on  $X$  has the **covering property** if  $I^{ext} = I^{int}$  for  $\Sigma_1^1$  sets, i.e. a  $\Sigma_1^1$  set  $A$  is in  $I^{int}$  iff  $A$  is in  $I^{ext}$ . This notions was introduced by Kechris in [4]. This is a quite strong property, in fact the only known ideals that have the covering property are  $K_\omega(X)$  and  $U_0$ . For  $K_\omega(X)$ , the classical perfect set theorem for  $\Sigma_1^1$  sets is the assertion that  $K_\omega(X)$  has the covering property. And for  $U_0$  is a theorem of Debs and Saint Raymond (see [2]).

In this article we undertake a study of this property from the descriptive set theoretic point of view. We will use as a paradigm the  $\sigma$ -ideal of countable closed sets, specifically the following five properties:

- (1) The classical perfect set theorem.
- (2) The collection of  $\Sigma_1^1$  countable sets is  $\Pi_1^1$  on the codes.
- (3) The effective version of the perfect set theorem says that a  $\Sigma_1^1$  countable set contains only hyperarithmetic points.
- (4) There is a largest  $\Pi_1^1$  set without perfect subset.
- (5) The perfect set theorem can be extended to  $\Sigma_2^1$  sets from large cardinals axioms. And it is false for  $\Pi_1^1$  sets in the constructible universe.

This article is divided into five sections respectively dealing with the five properties mentioned above. In fact we will show that similar results hold for  $\sigma$ -ideals of compact sets with the covering property.

## 2 The covering property and some related notions

We will work with the effective methods of descriptive set theory, so we assume that  $X$  is recursively presented (i.e. its metric is effective, see [9]). The collection of compact subsets of  $X$  becomes itself a compact, metric space under the usual metric:

$$\rho(K, L) = \begin{cases} \sup\{\max\{d(x, K), d(y, L)\} : x \in K, y \in L\} & , \text{ if } K, L \neq \emptyset \\ \text{diam}(X) & , \text{ if } K \text{ or } L = \emptyset \\ 0 & , \text{ if } K = L = \emptyset. \end{cases}$$

All topological and descriptive set theoretic notions concerning  $\mathcal{K}(X)$  refer to this space (for more details about the topology over  $\mathcal{K}(X)$  see [8] and the references given there). For instance, most of the times we will impose a definability condition over  $I$ , namely, it has to be a  $\Pi_1^1$  subset of  $\mathcal{K}(X)$ . We will use standard notions of descriptive set theory as in Moschovakis' book [9] and the notations from [8].

As we said in the introduction with each ideal  $I$  of closed subsets of  $X$ , there are two classes of (arbitrary) subsets of  $X$  associated with  $I$ :  $\mathbf{I}^{\text{int}}$  and  $\mathbf{I}^{\text{ext}}$ . The following concept was introduced by Kechris ([4]):

**Definition 2.1.** *We say that  $I$  has the **covering property**, if for every  $\Sigma_1^1$  set  $A \in I^{\text{int}}$ , there is a countable collection  $\{F_n\}$  of closed sets in  $I$  such that  $A \subseteq \bigcup_n F_n$ . And in general for a pointclass  $\Gamma$  we say that  $I$  has the covering property for  $\Gamma$ -sets, if for every set  $A \in \Gamma$  with  $A \in I^{\text{int}}$  there is a countable collection  $\{F_n\}$  of closed sets in  $I$  such that  $A \subseteq \bigcup_n F_n$ .*

Let us observe that for a  $\sigma$ -ideal  $I$  consisting of meager sets, the covering property implies that  $\Sigma_1^1$  sets in  $I^{\text{int}}$  are of first category, i.e., they are also small in the sense of category.

As we have mentioned before the classical perfect set theorem for  $\Sigma_1^1$  sets says that  $K_\omega(X)$  has the covering property. So, we can regard this property as an abstraction of the content of the perfect set theorem. Since in ZFC this theorem cannot be extended to  $\Pi_1^1$  sets, we do not expect to have (in ZFC) the covering property for  $\Pi_1^1$  sets (we will look at this problem in section 6).

In this section we will introduce some notions related with the covering property and show some structural and definability consequences of the covering property. As a corollary we will obtain a result of Kaufman about sets of extended uniqueness and also a partial answer to a question raised in [8].

The following notion is closely related to the covering property.

**Definition 2.2.** *An ideal  $I$  is **calibrated** if for every closed set  $F$  the following holds: If for some collection  $\{F_n\}$  of closed sets in  $I$ ,  $F - \bigcup_n F_n \in I^{\text{int}}$ , then  $F \in I$ .*

A typical calibrated ideal is the collection of closed null sets with respect to some Borel measure. On the other hand, the ideal of closed meager sets is not calibrated. Notice also that the covering property clearly implies calibration.

Let  $B$  be a hereditary subset of  $\mathcal{K}(X)$ .  $B_\sigma$  denotes the smallest  $\sigma$ -ideal (of closed sets) containing  $B$ , i.e.,  $K \in B_\sigma$  if there is a sequence  $\{K_n\}$  of elements of  $B$  such that  $K = \bigcup_n K_n$ . We say that  $I$  has a **Borel basis** if there is a Borel hereditary set  $B \subseteq I$  such that  $I = B_\sigma$ .  $I$  is called **locally non-Borel** if for every closed set  $F \notin I$ ,  $I \cap \mathcal{K}(F)$  is not Borel.

The only criterion known to show that an ideal has the covering property is the following theorem, which was originally used to show that the  $\sigma$ -ideal of closed set of uniqueness does not have a Borel basis (see [7] for a proof of both results).

**Theorem 2.3.** *(Debs-Saint Raymond [2]). Let  $I$  be a calibrated, locally non-Borel,  $\Pi_1^1$   $\sigma$ -ideal. If  $I$  has a Borel basis, then  $I$  has the covering property.*

□

Kechris [6] has asked the question of characterizing the  $\sigma$ -ideals which have the covering property. As we said, it implies calibration, but it is not known if the other hypotheses of the previous

theorem are necessary. Let us recall here that a  $\Pi_1^1$   $\sigma$ -ideal  $I$  satisfies the so called dichotomy theorem: It is either a true  $\Pi_1^1$  set or a  $G_\delta$  set (see [8]).

The usual way to show that the covering property fails for a  $\sigma$ -ideal  $I$  consisting of meager sets is by finding a dense  $G_\delta$  set  $G$  with  $G \in I^{int}$ . In fact, let us suppose such a  $G$  can be covered by a collection  $\{F_n\}$  of sets in  $I$ . Then by the Baire category theorem there is an  $n$  and an open set  $V$  such that  $V \cap G \neq \emptyset$  and  $V \cap G \subseteq F_n$ . As  $G$  is dense, we get  $V \subseteq \overline{V \cap G} \subseteq F_n$ , which contradicts that  $F_n$  is meager. In other words, the covering property fails for a  $G_\delta$  set. This is the case, for instance, when  $I$  consists of the null sets with respect to a Borel measure.

The following notion is quite useful: A non-empty set  $A$  is said to be **locally not in  $I$**  (or  **$I$ -perfect**), if for every open set  $V$  with  $V \cap A \neq \emptyset$ , we have that  $\overline{V \cap A} \notin I$ . Notice that  $A$  is  $I$ -perfect iff  $\overline{A}$  is  $I$ -perfect. Given a closed set  $F \notin I$ , there is a closed  $F' \subseteq F$  such that  $F'$  is locally not in  $I$ . In fact, let  $O = \bigcup \{V \subseteq X : V \text{ is open and } F \cap V \in I^{ext}\}$ . Put  $F' = F - O$ . It is easy to check that  $F'$  is locally not in  $I$ .  $F'$  is called the  $I$ -perfect kernel of  $F$ .

We will see later on that it is convenient to restrict attention to the covering property for  $\Pi_2^0$  sets. We have the following useful characterization of this notion

**Lemma 2.4.** *Let  $I$  be a  $\sigma$ -ideal of compact sets. The following are equivalent:*

- (i)  *$I$  has the covering property for  $\Pi_2^0$  sets.*
- (ii) *For each  $\Pi_2^0$  set  $G$  such that  $\overline{G}$  is locally not in  $I$ , we have  $G \notin I^{int}$ .*

**Proof:** (i) $\Rightarrow$ (ii). Let  $G$  be a  $G_\delta$  set such that  $M = \overline{G}$  is locally not in  $I$ . Suppose, towards a contradiction, that  $G \in I^{int}$ . By (i) there is a sequence  $\{F_n\}$  of sets in  $I$  such that  $G \subseteq \bigcup_n F_n$ . By the Baire category theorem there is an  $n$  and an open set  $V$  such that  $\emptyset \neq G \cap V \subseteq F_n$ . Hence  $\overline{V \cap G} = \overline{V \cap F_n} \subseteq F_n$ . So,  $\overline{V \cap G} \in I$ , which contradicts that  $M$  is locally not in  $I$ .

(ii) $\Rightarrow$ (i). Let  $G$  be a  $\Pi_2^0$  set in  $I^{int}$ . Assume towards a contradiction that  $G \notin I^{ext}$ . Let  $O = \bigcup \{V \subseteq X : V \text{ is an open set and } V \cap G \in I^{ext}\}$ . Let  $G' = G - O$ . As  $G \notin I^{ext}$ , then  $G' \neq \emptyset$ . It is clear that for all  $V$  open, if  $V \cap G' \neq \emptyset$  then  $V \cap G' \notin I^{ext}$ . Clearly  $G'$  is a  $\Pi_2^0$  set in  $I^{int}$  and for every open set  $V$ , if  $V \cap G' \neq \emptyset$  then  $\overline{V \cap G'} \notin I$ . Therefore  $M = \overline{G}$  is locally not in  $I$ , which contradicts (ii). □

**Remark:** A analogous result can be proved for the covering property for Borel sets as follows: Consider the  $\Delta_1^1$  topology  $\tau$ , i.e. the topology generated by the  $\Delta_1^1$  sets. This is a Polish topology. We say that a set  $A$  is  $\tau$ -locally not in  $I$  if for every  $\tau$ -open set  $V$  with  $V \cap A \neq \emptyset$  we have that  $\overline{V \cap A} \notin I$ . Then, as before, we have that the following are equivalent:

- (i)  $I$  has the covering property for  $\tau - G_\delta$  sets.
- (ii) For each  $\tau - G_\delta$  set  $G$  such that  $G$  is locally not in  $I$ , we have  $G \notin I^{int}$ .

The next type of ideals that we are going to consider are the thin ideals. This notion was introduced in [8] and it corresponds dually to the countable chain condition. We say that  $I$  is **thin** if every collection of pairwise disjoint closed sets not in  $I$  is at most countable. The typical example of thin ideal is the collection of null sets for some Borel measure. The next theorem relates thinness with the covering property.

**Theorem 2.5.** *Let  $I$  be a  $\sigma$ -ideal of closed sets which satisfies one of the following non triviality conditions:*

- (i)  $I \neq \mathcal{K}(X)$  and for every  $x \in X$ ,  $\{x\} \in I$ .
- (ii) Every  $K \in I$  is a meager set.

If  $I$  is thin, then  $I$  does not have the covering property for  $\Pi_2^0$  sets. Actually, if  $I$  is thin and (ii) holds, then there is a dense  $G_\delta$  set in  $I^{int}$ .

**Proof:** Assume first that (i) holds. Let  $O = \bigcup \{V \subseteq X : V \text{ is open and } V \in I^{ext}\}$ .  $O$  is the largest open set in  $I^{ext}$ . Put  $K = X - O$ ,  $K$  is locally not in  $I$  (if  $V \cap K \neq \emptyset$ , then  $\overline{V \cap K} \notin I$ , otherwise  $V \subseteq O$ ). As  $I \neq \mathcal{K}(X)$  and every singleton is in  $I$ , then  $K$  is a (non-empty) perfect set. Let  $G$  be a dense  $G_\delta$  subset of  $K$  with empty interior with respect to the relative topology of  $K$ . Let  $\{K_n\}$  be a maximal collection of pairwise disjoint closed subsets of  $G$  with each  $K_n \notin I$ . Each  $K_n$  is meager in  $K$ . Put  $F = \bigcup_n K_n$  and  $H = G - F$ . Then  $H$  is a dense (in  $K$ )  $G_\delta$  subset of  $K$ . Clearly  $H \in I^{int}$ , hence by 2.4  $I$  does not have the covering property for  $\Pi_2^0$  sets.

Now if (ii) holds, then  $X$  is locally not in  $I$ , hence the same proof applies. Finally, let's observe that in this case we get a dense  $G_\delta$  set in  $I^{int}$ . □

**Remark:** (i) Besides  $I \neq \mathcal{K}(X)$ , some other non-triviality condition has to be imposed on  $I$  in order to get the conclusion of 2.5, as the following example shows: let  $F \subseteq X$  be a countable closed set and  $V = X - F$ . Put  $I = \mathcal{K}(V)$ .  $I$  is thin, because  $K \notin I$  iff  $K \cap F \neq \emptyset$ . Thus there are only countable many disjoint sets not in  $I$ . However,  $I$  trivially satisfies the covering property (because  $V \in I^{ext}$  and if  $H \in I^{int}$  then  $H \subseteq V$ ).

(ii) We will use 2.5 usually as follows. Suppose that every Borel set in  $I^{int}$  is of the first category ( $\Pi_2^0$  sets suffice). Then  $I$  is not thin. Just notice that in this case every set in  $I$  is meager.

The following notion was introduced in [8]. A set  $A \subseteq X$  is called **I-thin** if there is no uncountable family of pairwise disjoint closed subsets of  $A$  which are not in  $I$ . In other words,  $A$  is  $I$ -thin if the restriction of  $I$  to  $\mathcal{K}(A)$  is a thin ideal. Given an ideal  $I$  define another ideal  $J_I$  as follows:

$$K \in J_I \text{ iff } K \text{ is } I\text{-thin.}$$

It was proved in [8] that if  $I$  is a  $\Pi_1^1$  calibrated  $\sigma$ -ideal then so is  $J_I$ . It was asked there to find out for a given  $I$  whether  $J_I = I$ . In relation with this question we have the following

**Corollary 2.6.** *Let  $I$  be a  $\sigma$ -ideal of closed subsets of  $X$  containing all singletons. If  $I$  has the covering property for  $\Pi_2^0$  sets, then  $I = J_I$ .*

**Proof:** It is clear that  $I \subseteq J_I$ . Now, let  $F$  be a closed set not in  $I$ . We want to show that  $F \notin J_I$ . We can assume without loss of generality that  $F$  is locally not in  $I$ . Hence as  $I$  contains all singletons,  $F$  is perfect. Put  $\tilde{I} = \mathcal{K}(F) \cap I$ .  $\tilde{I}$  is non trivial in the sense of 2.5 (i) and it has the covering property for  $\Pi_2^0$  sets: if  $H \subseteq F$  is a  $\Pi_2^0$  set in  $\tilde{I}^{int}$  then  $H \in I^{int}$ . Hence, by the covering property for  $I$ ,  $H \in I^{ext}$ . This clearly implies that  $H \in \tilde{I}^{ext}$ . Therefore, by 2.5  $\tilde{I}$  is not thin, i.e.,  $F \notin J_I$ . □

**Corollary 2.7.** *(Kaufman) Let  $U_0$  denote the  $\sigma$ -ideal of closed set of extended uniqueness in the unit circle. Then  $U_0 = J_{U_0}$ .*

**Proof:** Debs and Saint Raymond [2] have shown that  $U_0$  has the covering property. □

Theorem 2.5 says that a non trivial  $\Pi_1^1$  thin  $\sigma$ -ideal  $I$  does not have the covering property. In [8] it was asked whether for an  $I$  that was also calibrated we have that  $I$  has to be  $\Pi_2^0$ . The next theorem is a partial answer to this question.

**Theorem 2.8.** *If  $I$  is a calibrated, thin,  $\Pi_1^1$   $\sigma$ -ideal of closed sets with a Borel basis, then  $I$  is  $\Pi_2^0$ .*

**Proof:** Let  $\{F_n\}$  be a maximal pairwise disjoint countable collection of closed sets such that for each  $n$ ,  $F_n \notin I$  and  $I \cap \mathcal{K}(F_n)$  is  $\Pi_2^0$ . Put  $F = \bigcup_n F_n$  and  $H = X - F$ . We claim that  $H \in I^{int}$ . Granting this claim we have:

$$K \in I \text{ iff } (\forall n)(K \cap F_n \in I). \quad (*)$$

The direction  $(\Rightarrow)$  is trivial. On the other hand, let  $K \subseteq X$  be a closed set. Then  $K = (K \cap H) \cup \bigcup_n (K \cap F_n)$ . Suppose that each  $K \cap F_n \in I$ . As  $I$  is calibrated and  $K \cap H \in I^{int}$ , then  $K \in I$ .

Now, the map  $K \mapsto K \cap F_n$  is Borel, so  $(*)$  says that  $I$  is Borel. Therefore by the Dichotomy theorem (see [8] theorem 1.7)  $I$  is  $\Pi_2^0$ .

It remains to show that  $H$  is in  $I^{int}$ . Suppose not towards a contradiction. Let  $M \subseteq H$  be a closed set locally not in  $I$ . Since  $\{F_n\}$  is maximal then for every  $x \in M$ ,  $\{x\} \in I$ . Hence  $M$  is a perfect set. Consider the  $\sigma$ -ideal  $I_0 = \mathcal{K}(M) \cap I$ .  $I_0$  is clearly a calibrated, thin (non-trivial as in 2.5)  $\Pi_1^1$   $\sigma$ -ideal with a Borel basis. As  $\{F_n\}$  is maximal, for every  $F \subseteq M$  with  $F \notin I_0$  we have that  $\mathcal{K}(F) \cap I_0 = \mathcal{K}(F) \cap I$  is not  $\Pi_2^0$ . Hence  $I_0$  is locally non Borel and thus all the hypotheses of the Debs-Saint Raymond theorem (2.3) are satisfied. Therefore  $I_0$  has the covering property, but also it is non trivial and thin which contradicts 2.5. □

This raises the following question: Does every calibrated, thin  $\Pi_1^1$   $\sigma$ -ideal have a Borel basis ?

### 3 Complexity of the codes

As we said in the introduction, another feature of the  $\sigma$ -ideal of countable sets is that it is  $\Pi_1^1$  on the codes of  $\Sigma_1^1$  sets. We will present an abstract version of this result as a consequence of the covering property. The key notion involucred is the following

**Definition 3.1.** *An ideal  $I$  is **strongly calibrated** if for every closed set  $F \subseteq X$  with  $F \notin I$  and every  $\Pi_2^0$  set  $H \subseteq X \times 2^\omega$  such that  $proj(H) = F$ , there is a closed set  $K \subseteq H$  such that  $proj(K) \notin I$ .*

This notion was introduced in [8]. It resembles the conclusion of Choquet's capacitability theorem and in fact this theorem implies that the  $\sigma$ -ideal of closed measure zero sets for a collection of Borel measures is strongly calibrated: Let  $\mathcal{M}$  be a collection of Borel measures on  $X$  and let  $I = Null(\mathcal{M})$ . Let  $Q \subseteq X \times 2^\omega$  be a  $\Pi_2^0$  set such that  $proj(Q) = F \notin I$ , and say  $\mu(F) > 0$  for some  $\mu \in \mathcal{M}$ . Define a capacity  $\gamma$  on  $X \times 2^\omega$  as follows:

$$\gamma(A) = \mu^*(proj(A)), \text{ for } A \subseteq X \times 2^\omega.$$

As  $Q$  is  $\Pi_2^0$  and  $\gamma(Q) > 0$ , by Choquet's capacitability theorem there is a compact set  $K \subseteq Q$  such that  $\gamma(K) > 0$ . Hence  $proj(K) \notin I$ .

These type of ideals have the property that the collection of  $\Sigma_1^1$  sets in  $I^{int}$  is  $\Pi_1^1$  on the codes of  $\Sigma_1^1$  sets (assuming that  $I$  is  $\Pi_1^1$ ). The usual argument to show this uses the capacitability theorem. We show next that strongly calibrated  $\sigma$ -ideals also have this property.

**Theorem 3.2.** *Let  $I$  be a  $\Pi_1^1$  strongly calibrated  $\sigma$ -ideal of closed subsets of  $X$ . Then the collection of  $\Sigma_1^1$  sets in  $I^{int}$  is  $\Pi_1^1$  in the codes of  $\Sigma_1^1$  sets.*

**Proof:** Let  $\mathcal{U} \subseteq 2^\omega \times X$  be a  $\Sigma_1^1$  universal set for  $\Sigma_1^1$  subsets of  $X$ . Let  $Q \subseteq (2^\omega \times X) \times 2^\omega$  be a  $\Pi_2^0$  set such that  $\mathcal{U} = \text{proj}(Q)$ . Consider the following relation

$$R(F, \alpha) \text{ iff } F \subseteq \mathcal{U}_\alpha \text{ \& } F \notin I.$$

Then we have

$$\mathcal{U}_\alpha \notin I^{int} \text{ iff } (\exists F) R(F, \alpha).$$

Hence it suffices to show that  $R$  is  $\Sigma_1^1$ . We claim that

$$R(F, \alpha) \text{ iff } (\exists K \in \mathcal{K}(2^\omega \times X))(K \subseteq Q^\alpha \text{ \& } \text{proj}(K) \notin I). \quad (*)$$

The direction  $\Leftarrow$  clearly holds. For the other, suppose that  $R(F, \alpha)$  holds and put  $H = Q^\alpha \cap (2^\omega \times F)$ . Then  $\text{proj}(H) = F$ . As  $H$  is  $\Pi_2^0$ , by strong calibration there is a closed  $K \subseteq H$  such that  $\text{proj}(K) \notin I$ , this set  $K$  clearly works.

To see that  $(*)$  is  $\Sigma_1^1$  we use the uniformization theorem for relations with  $K_\sigma$  sections (see theorem 4F.16 in [9]) and we get that  $K \subseteq Q^\alpha$  is  $\Delta_1^1$ . (In fact, since we are working in compact spaces the projection of a  $\Sigma_2^0$  is also  $\Sigma_2^0$ , with this in mind it is easy to check that  $K \subseteq Q^\alpha$  is a  $\Pi_2^0$  relation on  $K$  and  $\alpha$ ). On the other hand, by a similar argument it is easy to see that the function  $K \mapsto \text{proj}(K)$  is  $\Delta_1^1$ -recursive (it is clearly continuous).

□

Strong calibration implies calibration (see [8] page 283). Also, one can take projections of  $\Sigma_1^1$  subsets of any compact Polish space in the definition of strong calibration as the following proposition shows. This sometimes makes this notion easier to use.

**Lemma 3.3.** *Strong calibration is equivalent to any of the following statements.*

- (i) *If  $F \subseteq X$  is a closed set not in  $I$  and  $Q \subseteq X \times 2^\omega$  is a  $\Sigma_1^1$  set such that  $\text{proj}(Q) = F$ , then there is a closed set  $K \subseteq Q$  such that  $\text{proj}(K) \notin I$ .*
- (ii) *Let  $Y$  be a compact Polish space. If  $F \subseteq X$  is a closed set not in  $I$  and  $Q \subseteq X \times Y$  is a  $\Sigma_1^1$  set such that  $\text{proj}(Q) = F$ , then there is a closed set  $K \subseteq Q$  such that  $\text{proj}(K) \notin I$ .*

**Proof:** (ii) follows from (i) because for any compact Polish space  $Y$  there is a continuous surjection  $f : 2^\omega \rightarrow Y$ .

To prove (i), let  $Q \subseteq X \times 2^\omega$  be a  $\Sigma_1^1$  set as in the hypothesis of (i). Let  $P \subseteq X \times 2^\omega \times 2^\omega$  be a  $\Pi_2^0$  set such that  $\text{proj}(P) = Q$ . Let  $f : 2^\omega \rightarrow 2^\omega \times 2^\omega$  be an homeomorphism, say  $f = (f_0, f_1)$ . Define  $P^* \subseteq X \times 2^\omega$  by

$$(x, \alpha) \in P^* \text{ iff } (x, f_0(\alpha), f_1(\alpha)) \in P.$$

Then  $P^*$  is  $\Pi_2^0$  and clearly  $\text{proj}(P^*) = F$ . So by strong calibration, there is a closed  $K^* \subseteq P^*$  such that  $\text{proj}(K^*) \notin I$ . Define  $K \subseteq X \times 2^\omega$  by  $(x, \alpha) \in K$  iff  $(\exists \beta)((x, f^{-1}(\alpha, \beta))) \in K^*$ . It is easy to check that  $K$  is a closed subset of  $Q$  and  $\text{proj}(K) = \text{proj}(K^*)$ .

□

As we said before we have the following

**Theorem 3.4.** *Let  $I$  be a  $\sigma$ -ideal of closed subsets of  $X$ . If  $I$  has the covering property for  $\Pi_2^0$  sets, then  $I$  is strongly calibrated.*

**Proof:** Let  $F$  be a closed set not in  $I$  and  $Q \subseteq X \times 2^\omega$  be a  $\Pi_2^0$  set such that  $F = \text{proj}(Q)$ . Without loss of generality we can assume that  $F$  is locally not in  $I$ . By the von Neumann selection theorem (see 4E.9 in [9]) there is a Baire measurable function  $f$  such that for all  $x \in F, (x, f(x)) \in Q$ . By the analog of the Lusin's theorem for category (see [10]), there is a  $G_\delta$  set  $G \subseteq F$  dense in  $F$ , such that  $f$  is continuous on  $G$ . Since  $I$  has the covering property for  $\Pi_2^0$  sets, then by 2.4,  $G \notin I^{int}$ . Thus, there is a closed set  $K \subseteq F$  with  $K \notin I$ . Let  $K^* = \text{graph of } f \text{ restricted to } K$ . As  $f$  is continuous on  $K$ , then  $K^*$  is a closed set and clearly  $\text{proj}(K^*) = K$ . This finishes the proof.  $\square$

**Corollary 3.5.** *Let  $I$  be a  $\Pi_1^1$  locally non Borel  $\sigma$ -ideal with a Borel basis. Then  $I$  is calibrated iff  $I$  is strongly calibrated.*

**Proof:** It was proved in [8](page 283) that strong calibration implies calibration. On the other hand, by the Debs-Saint Raymond theorem (2.3) every  $\sigma$ -ideal as in the hypothesis above has the covering property. Hence, by previous theorem it is strongly calibrated.  $\square$

From the proof of 3.4 one gets the following: Let's say that an ideal  $I$  has the *continuity property* if for every Baire measurable function  $f$  with  $\text{dom}(f) = F \notin I$  ( $F$  a closed set), there is a closed set  $K \subseteq F, K \notin I$  and  $f$  continuous on  $K$ .

**Corollary 3.6.** *(of the proof of 3.4) Let  $I$  be a  $\sigma$ -ideal of closed subsets of  $X$ .*

- (i) *If  $I$  has the covering property for  $\Pi_2^0$  sets, then  $I$  has the continuity property.*
- (ii) *If  $I$  has the continuity property, then  $I$  is strongly calibrated.*

$\square$

**Remark:** Observe that if  $I$  is strongly calibrated, then  $I$  has the continuity property for Borel functions: Just apply the definition of strong calibration to the graph of  $f$ .

Strong calibration is not equivalent to the covering property for  $\Pi_2^0$  sets, because as we have already mentioned  $\text{Null}(\mu)$  is strongly calibrated but it does not have the covering property.

Calibration is equivalent to saying that  $I^{int} \cap \Pi_2^0(X)$  is a  $\sigma$ -ideal (see Proposition 1 §3 in [8]). The next lemma shows that for strong calibration we get a similar result for  $\Sigma_1^1$  sets.

**Lemma 3.7.** *Let  $I$  be a strongly calibrated  $\sigma$ -ideal. Then*

- (i) *If  $F$  is a closed set such that  $F = P \cup \bigcup_n F_n$ , for some  $\Sigma_1^1$  set  $P$  in  $I^{int}$  and each  $F_n$  in  $I$ , then  $F \in I$ . In particular  $I$  is calibrated.*
- (ii)  *$\{P \subseteq X : P \text{ is a } \Sigma_1^1 \text{ set in } I^{int}\}$  is a  $\sigma$ -ideal.*
- (iii) *Define a collection  $J \subseteq \mathcal{K}(X \times 2^\omega)$  as follows:*

$$K \in J \text{ iff } \text{proj}(K) \in I$$

*Then  $J$  is a calibrated  $\sigma$ -ideal.*

**Proof:** (i) Let  $F = P \cup \bigcup_n F_n$  be a closed set not in  $I$  with  $P$  a  $\Sigma_1^1$  set and each  $F_n$  in  $I$ . We will show that  $P \notin I^{int}$ . Let  $G \subseteq X \times 2^\omega$  be a  $\Pi_2^0$  set such that  $\text{proj}(G) = P$ . Put

$$Q = (G \times \{0\}) \cup \bigcup_n (F_n \times 2^\omega \times \{1\})$$

$Q \subseteq X \times (2^\omega \times (\omega + 1))$  and  $\text{proj}(Q) = F$ . By strong calibration there is  $K \subseteq Q$  closed such that  $\text{proj}(K) \notin I$ . Now, we have

$$K = K \cap (G \times \{0\}) \cup \bigcup_n K \cap (F_n \times 2^\omega \times \{1\}).$$

Hence

$$\text{proj}(K) = \text{proj}(K \cap (G \times \{0\})) \cup \bigcup_n \text{proj}(K \cap (F_n \times 2^\omega \times \{1\})).$$

Since  $K \cap (G \times \{0\})$  is closed in  $X \times (2^\omega \times (\omega + 1))$  and  $\text{proj}(K \cap (F_n \times 2^\omega \times \{1\})) \subseteq F_n \in I$ , then  $\text{proj}(K \cap (G \times \{0\})) \notin I$ . Thus  $\text{proj}(G) = P \notin I^{int}$ .

We show (iii) first. It is clear that  $J$  is a  $\sigma$ -ideal. Let  $K = G \cup \bigcup_n K_n$ , where  $K \subseteq X \times 2^\omega$  is closed,  $G$  is a  $\Pi_2^0$  set in  $J^{int}$  and each  $K_n$  is in  $J$ . Now,  $\text{proj}(K) = \text{proj}(G) \cup \bigcup_n \text{proj}(K_n)$ . As  $\text{proj}(K_n)$  is a closed set in  $I$ , it suffices to show that  $\text{proj}(G) \in I^{int}$  and then apply (i). Let  $F \subseteq \text{proj}(G)$  and suppose toward a contradiction that  $F \notin I$ . By strong calibration there is  $K \subseteq (F \times 2^\omega) \cap G$  closed such that  $\text{proj}(K) \notin I$ . This contradicts that  $G$  in  $J^{int}$ .

(ii) It is easy to check (as in (iii)) that strong calibration implies that

$$\{P \subseteq X : P \in \Sigma_1^1(X) \cap I^{int}\} = \{\text{proj}(G) : G \in \Pi_2^0(X \times 2^\omega) \cap J^{int}\}.$$

Since  $J$  is calibrated the collection of  $\Pi_2^0$  sets in  $J^{int}$  is a  $\sigma$ -ideal (in fact  $\Sigma_3^0$  sets, see Proposition 1§3 in [8]), from which the claim follows.  $\square$

The next lemma relates the covering property of  $I$  and  $J$ , it will be used in section 5.

**Lemma 3.8.** *Let  $I$  be a  $\sigma$ -ideal and  $J$  be the  $\sigma$ -ideal defined in 3.6 (iii). Then the following are equivalent:*

- (i)  $J$  has the covering property.
- (ii)  $J$  has the covering property for  $\Pi_2^0$  sets.
- (iii)  $I$  has the covering property.

**Proof:** Clearly (i)  $\Rightarrow$  (ii).

(ii)  $\Rightarrow$  (iii). Let  $P$  be a  $\Sigma_1^1$  set in  $I^{int}$  and  $G \subseteq X \times 2^\omega$  be a  $\Pi_2^0$  set such that  $\text{proj}(G) = P$ . Clearly  $G \in J^{int}$ . Hence there are closed sets  $K_n \in J$  such that  $G \subseteq \bigcup_n K_n$ . Each  $\text{proj}(K_n) \in I$  and  $\text{proj}(G) \subseteq \bigcup_n \text{proj}(K_n)$ .

(iii)  $\Rightarrow$  (i). Let  $G \subseteq X \times 2^\omega$  be a  $\Sigma_1^1$  set with  $G \in J^{int}$ . By 3.4  $I$  is strongly calibrated, hence (as in the proof of (ii) in 3.7)  $\text{proj}(G) \in I^{int}$ . So, there are closed sets  $F_n$  in  $I$  such that  $\text{proj}(G) \subseteq \bigcup_n F_n$ . Thus  $G \subseteq \bigcup_n F_n \times 2^\omega$  and clearly for all  $n$ ,  $F_n \times 2^\omega \in J$ .  $\square$

If  $I$  has the covering property then for every  $\Sigma_1^1$  set  $A \in I^{int}$  there is a Borel (actually an  $F_\sigma$ ) set  $B \in I^{int}$  with  $A \subseteq B$ . The next result shows that this is also a consequence of strong calibration, which in particular says that the covering property for Borel sets implies the covering property (for  $\Sigma_1^1$  sets).

**Theorem 3.9.** *Let  $I$  be a strongly calibrated  $\Pi_1^1$   $\sigma$ -ideal. Let  $A$  be a  $\Sigma_1^1$  set in  $I^{int}$ . Then there is a  $\Delta_1^1$  set  $B \in I^{int}$  such that  $A \subseteq B$ . Therefore, if we let*

$$H(I) = \bigcup \{B \subseteq X : B \text{ is } \Delta_1^1 \text{ and } B \in I^{int}\},$$

we have

- (i)  $H(I)$  is a  $\Pi_1^1$  set in  $I^{int}$ .
- (ii) For every  $\Sigma_1^1$  set  $A$ ,  $A \in I^{int}$  iff  $A \subseteq H(I)$ .

**Proof:** The first claim follows from the reflection principle but we give a direct proof anyway. Let  $A$  be a  $\Sigma_1^1$  set in  $I^{int}$  and put  $P = X - A$ . Let  $\varphi$  be a  $\Pi_1^1$  norm on  $P$  and consider

$$M = \{x \in X : \{y : \neg(y <_\varphi^* x)\} \in I^{int}\}.$$

As in the proof of proposition 3.2 we have that  $M$  is  $\Pi_1^1$ . We claim that  $A \subseteq M$ . In fact, if  $x \in A$  then by definition of  $<_\varphi^*$  we have that

$$\{y : \neg(y <_\varphi^* x)\} = A.$$

By separation, let  $B \subseteq M$  be a  $\Delta_1^1$  set with  $A \subseteq B \subseteq M$ . If  $A = B$  we are done. Else let  $\xi$  be the least ordinal in  $\{\varphi(x) : x \in B\}$  and let  $x \in B$  with  $\varphi(x) = \xi$ . Then

$$B \subseteq \{y : \neg(y <_\varphi^* x)\}.$$

Hence  $B \in I^{int}$ .

From the proposition 3.7 we know that the collection of  $\Sigma_1^1$  sets in  $I^{int}$  form a  $\sigma$ -ideal, so  $H(I) \in I^{int}$ . As in the proof of 3.2 we can show that  $H(I)$  is  $\Pi_1^1$ . This proves (i). And (ii) follows from (i) and the first claim. □

The set  $H(I)$  can be thought as an abstract version of the hyperarithmetic reals. A better description of it will be given in the next section. By Theorem 3.4 the covering property for  $G_\delta$  sets implies strong calibration, thus from the relativized version of the previous theorem we immediately get

**Theorem 3.10.** *Let  $I$  be a  $\Pi_1^1$   $\sigma$ -ideal. If  $I$  has the covering property for Borel sets, then it has the covering property.* □

## 4 Analog of the hyperarithmetic reals

One of the consequence of the effective perfect set theorem is that a countable  $\Sigma_1^1$  set contains only hyperarithmetic points, i.e.,  $\Delta_1^1$  points. We will present an abstract version of this result for  $\Pi_1^1$   $\sigma$ -ideals with the covering property. The main theorem is the following strengthening of 3.9

**Theorem 4.1.** *Let  $I$  be a  $\Pi_1^1$   $\sigma$ -ideal on  $2^\omega$  with the covering property. Let*

$$H(I) = \bigcup \{[T] : T \text{ is } \Delta_1^1 \text{ binary tree and } [T] \in I\},$$

we have

- (i)  $H(I)$  is a  $\Pi_1^1$  set in  $I^{ext}$ .
- (ii) For every  $\Sigma_1^1$  set  $A$ ,  $A \in I^{int}$  iff  $A \subseteq H(I)$ .

The key lemma used in the proof of this theorem is the following result due to Barua-Srivatsa ([1]), its proof is similar to the proof of theorem 3.9 and it is a bit different to the one given in [1].

**Lemma 4.2.** (see Barua-Srivatsa [1]) Let  $I$  be a  $\Pi_1^1$   $\sigma$ -ideal on  $2^\omega$  and  $T$  a  $\Sigma_1^1$  binary tree such that  $[T] \in I$ . Then there is a  $\Delta_1^1$  binary tree  $S$  such that  $[T] \subseteq [S]$  and  $[S] \in I$

**Proof:** Let  $Seq$  denote the collection of binary sequences. Fix a  $\Pi_1^1$  norm  $\varphi$  on  $Seq - T$  and consider the following sets: Let

$$A_s = \{t \in Seq : \exists t' \text{ extending } t \ \& \ \neg(t' <_\varphi^* s)\}$$

Notice that  $A_s$  is a tree. Let

$$M = \{s \in Seq : [A_s] \in I\}$$

As the set  $A_s$  is  $\Sigma_1^1$  one can easily check (as in 3.2) that the property  $[A_s] \in I$  is  $\Pi_1^1$ . Hence  $M$  is a  $\Pi_1^1$  set. Let us observe that  $T \subseteq M$ : If  $s \in T$ , it is easy to see that  $A_s = T$  hence  $s \in M$ . So, unless  $T$  is  $\Delta_1^1$  (in which case there is nothing to prove), there is  $s \in M - T$ . Thus we let

$$\alpha = \text{least ordinal of } \{\varphi(s) : s \in M\}$$

and let  $s_0 \in M$  such that  $\varphi(s_0) = \alpha$ .

It is clear that  $M \subseteq A_{s_0}$ : If  $s \in M$  then  $\neg(s <_\varphi^* s_0)$ , hence  $s \in A_{s_0}$ . Let  $B$  be a  $\Delta_1^1$  set such that  $T \subseteq B \subseteq M$  and let  $S = \{t \in Seq : \exists t' \text{ extending } t \ \& \ t' \in B\}$ .  $S$  is a  $\Delta_1^1$  tree and thus  $S \subseteq A_{s_0}$ . Hence  $[S] \in I$ . □

Now we are ready to give

**Proof of theorem 4.1 :** (i) It is obvious that  $H(I)$  is in  $I^{ext}$ . And by the theorem on restricted quantification (4D.3 in [9]) we get that  $H(I)$  is  $\Pi_1^1$ .

To see (ii), let  $H = H(I)$ , and suppose  $A \not\subseteq H$ , towards a contradiction. Let  $A^* = A - H$ .  $A^*$  is a  $\Sigma_1^1$  set in  $I^{int}$ . So, let  $\{K_n\}$  be closed sets in  $I$  such that  $A^* \subseteq \bigcup_n K_n$ . By working with the  $\Sigma_1^1$ -topology and by the Baire category theorem we know that there is a  $\Sigma_1^1$  set  $V$  such that  $\emptyset \neq V \cap A^* \subseteq K_n$  for some  $n$ . Let  $T$  be the tree of  $\overline{V \cap A^*}$ .  $T$  is clearly  $\Sigma_1^1$  and  $[T] \in I$ . Hence by lemma 4.2 there is a  $\Delta_1^1$  tree  $S$  such that  $T \subseteq S$  and  $[S] \in I$ . Thus  $[S] \subseteq H$ , which contradicts that  $A^* \cap H = \emptyset$ . This finishes the proof. □

As a corollary we get that the covering property holds effectively.

**Corollary 4.3.** (see Barua-Srivatsa [1]) Let  $I$  be a  $\Pi_1^1$   $\sigma$ -ideal on  $2^\omega$  with the covering property. Let  $A$  be a  $\Sigma_1^1$  set in  $I^{int}$ . Then there is a  $\Delta_1^1$  recursive function  $f : \mathbb{N} \rightarrow \omega^\omega$  such that for all  $n$   $f(n)$  is a binary tree with  $[f(n)] \in I$  and  $A \subseteq \bigcup_n [f(n)]$ .

**Proof:** The proof is an standard application of the selection principle.

By separation there is a  $\Delta_1^1$  set  $A^*$  such that  $A \subseteq A^* \subseteq H(I)$ . Let  $\mathbf{d}(i)$  be the canonical function that enumerates the  $\Delta_1^1$  points (see [9] 4D.2). Consider the following relation:

$$D(x, i) \text{ iff } \mathbf{d}(i) \downarrow \ \& \ \mathbf{d}(i) \text{ codes a } \Delta_1^1 \text{ binary tree } T \ \& \ x \in [T] \ \& \ [T] \in I$$

It is easy to check that  $D$  is  $\Pi_1^1$ .

We have that for all  $x \in A^*$ , there is  $i$  such that  $D(x, i)$  holds. Hence by the  $\Delta$ -selection principle (see [9] 4B.5), there is a  $\Delta_1^1$ -recursive function  $g : 2^\omega \rightarrow \omega$  such that for all  $x \in A^*$   $D(x, g(x))$  holds. Let  $R$  be the range of  $g$ ,  $R$  is  $\Sigma_1^1$ . Put

$$S = \{i : \mathbf{d}(i) \downarrow \text{ \& } \mathbf{d}(i) \text{ codes a } \Delta_1^1 \text{ binary tree } T \text{ with } [T] \in I\}.$$

Then  $S$  is  $\Pi_1^1$  and  $R \subseteq S$ . So, by separation there is a  $\Delta_1^1$  set  $R^*$  such that  $R \subseteq R^* \subseteq S$ . Define  $f$  as follows:

$$f(i) = \begin{cases} T & \text{if } i \in R^* \text{ and } \mathbf{d}(i) \text{ codes } T \\ \emptyset & \text{otherwise} \end{cases}$$

$f$  is clearly  $\Delta_1^1$ -recursive. For all  $i$ ,  $[f(i)] \in I$  and  $A \subseteq \bigcup_i f(i)$ . Also  $f(i)$  is a  $\Delta_1^1$  binary tree.  $\square$

## 5 The Largest $\Pi_1^1$ set in $I^{int}$

It is a well known fact that there is a largest  $\Pi_1^1$  thin set, i.e. a set without a perfect subset. This set is denoted by  $C_1$  and it is characterized by  $\alpha \in C_1$  iff  $\alpha \in L_{\omega_1^\alpha}$  (see [3] and [4] for similar results on  $\sigma$ -ideals on  $\omega^\omega$  defined by games). Another consequence of the covering property for a  $\sigma$ -ideal  $I$  is that there is a largest  $\Pi_1^1$  set in  $I^{int}$ . In this section we will present a proof of this fact. Moreover, for  $\sigma$ -ideals defined on  $2^\omega$  such a set can be characterized in a similar fashion as  $C_1$ .

There is a theorem due to Kechris (see [3] 1A-2) that gives sufficient conditions for the existence of such a largest  $\Pi_1^1$  set for  $\sigma$ -ideals of subsets of  $X$ . One of these conditions is the so called  $\Pi_1^1$ -additivity. We will show next that for every  $\sigma$ -ideal  $I$  of meager subsets of  $X$ , if  $I$  has the covering property, then  $I^{int}$  is  $\Pi_1^1$ -additive. The proof is based on a representation of  $I$  as the common meager closed sets for a collection of Polish topologies on  $X$ .

**Definition 5.1.** For every topology  $\tau$  on  $X$ , let  $Meager(\tau)$  be the collection of  $\tau$ -meager sets. We say that a topology  $\tau$  on  $X$  is **compatible with**  $I$  if  $\tau$  extends the original topology on  $X$ , every  $\tau$ -open set is Borel and  $I \subseteq Meager(\tau)$ .

Observe that in this case the Borel structure of  $X$  and  $(X, \tau)$  are the same. In particular every  $C$ -measurable subset  $B \subseteq X$  has the property of Baire with respect to  $\tau$  ( $C$  is the least  $\sigma$ -algebra containing the open sets and closed under the Suslin operation).

**Lemma 5.2.** Let  $I$  be a  $\sigma$ -ideal of meager closed subsets of a compact Polish space  $X$ . Then we have

$$I = \bigcap \{Meager(\tau) \cap \mathcal{K}(X) : \tau \text{ is a Polish topology on } X \text{ compatible with } I\}.$$

**Proof:** One direction is obvious. Let  $K \notin I$ . We want to find a Polish topology  $\tau$  on  $X$  compatible with  $I$  and such that  $K$  is not  $\tau$ -meager. Without loss of generality we assume that  $K$  is locally not in  $I$ . Let  $\tau_0$  be the given topology on  $X$  and consider the topology  $\tau$  generated by

$$\tau_0 \cup \{V \cap K : V \in \tau_0\}.$$

It is a standard fact that  $\tau$  is the least Polish topology for which  $K$  is  $\tau$ -clopen. It remains only to show that  $I \subseteq Meager(\tau)$ . But this is clear, because as  $K$  is locally not in  $I$ , for every  $V \in \tau_0$  such that  $V \cap K \neq \emptyset$  we have that  $\overline{V \cap K} \notin I$ . Hence for every  $F \in I$ ,  $V \cap K \not\subseteq F$ .  $\square$

Let us introduce the following

**Definition 5.3.** We say that a subset of  $X$  is  **$I$ -meager** if it is  $\tau$ -meager for every topology  $\tau$ -compatible with  $I$ .

Thus the previous lemma says that a closed set is in  $I$  iff is  $I$ -meager. As we said before the key fact in the proof of the existence of the largest  $\Pi_1^1$  set in  $I^{int}$  is the following

**Theorem 5.4.** *Let  $I$  be a  $\sigma$ -ideal of meager subsets of  $X$  with the covering property and let  $B$  be a subset of  $X$  with the property of Baire with respect to every Polish topology compatible with  $I$ . The following are equivalent:*

- (i)  $B \in I^{int}$ .
- (ii)  $B$  is  $I$ -meager.

**Proof:** (i)  $\Rightarrow$  (ii). Suppose that  $B$  is not  $\tau$ -meager for some topology  $\tau$  compatible with  $I$ . As  $B$  has the property of Baire for  $\tau$ , then there is a  $\tau$ -open set  $V$  such that  $B$  is  $\tau$ -comeager in  $V$ . So, let  $G$  be a  $\tau$ - $G_\delta$  set  $\tau$ -dense in  $V$  and  $G \subseteq B$ . As  $\tau$  consists of Borel sets then  $G$  is also Borel. We claim that  $G \notin I^{int}$ . Otherwise, as  $I$  has the covering property, there are closed sets  $\{F_n\}$  in  $I$  such that  $G \subseteq \bigcup_n F_n$ . Each  $F_n$  is  $\tau$ -closed, hence by the Baire category theorem there is a  $\tau$ -open set  $W$  and an integer  $n$  such that  $\emptyset \neq W \cap G \subseteq F_n$ . But as  $G$  is  $\tau$ -dense in  $V$  we get that  $F_n$  is not  $\tau$ -meager, which contradicts that  $\tau$  is compatible with  $I$ .

(ii)  $\Rightarrow$  (i). It follows immediately from the previous lemma.  $\square$

If we trace back how much the covering property is needed to prove this theorem we see that it would be sufficient with the covering property for  $G_\delta$  sets. This is because the topologies used in the proof of 5.2 admit a basis consisting of  $G_\delta$  sets in the original topology of  $X$ . In other words, the proof of 5.2 shows that

$$I = \bigcap \{ \text{Meager}(\tau_K) \cap \mathcal{K}(X) : K \text{ is a } I\text{-perfect closed set and } \tau_K \text{ is the canonical Polish topology for which } K \text{ is clopen} \}.$$

In fact the conclusion of the previous theorem is equivalent to the covering property for  $G_\delta$  sets, as we show next.

**Lemma 5.5.** *Let  $I$  be a  $\sigma$ -ideal of meager subsets of  $X$ .  $I$  has the covering property for  $G_\delta$  sets iff  $I\text{-meager} = I^{int}$  for sets with the Baire property with respect to every topology compatible with  $I$ .*

**Proof:** One direction follows from the previous theorem and the remark we did after it. For the other direction let  $G \subseteq X$  be a  $G_\delta$  set such that  $\overline{G}$  is  $I$ -perfect. We want to show that  $G \notin I^{int}$ . Suppose not, towards a contradiction. Thus by hypothesis  $G$  is  $I$ -meager. Let  $K = \overline{G}$  and  $\tau = \tau_K$  be the canonical topology for  $K$ . As  $\tau$  is compatible with  $I$  then  $G$  is  $\tau$ -meager. Let  $\{F_n\}$  be a collection of  $\tau$ -meager  $\tau$ -closed sets such that  $G \subseteq \bigcup_n F_n$ . By the usual argument with the Baire category theorem there is a  $\tau$ -basic open set  $V \cap K$  such that  $V \cap K \cap G = V \cap G \subseteq F_n$  for some  $n$ . And it is easy to check that this implies that  $F_n$  is not  $\tau$ -meager, which is a contradiction.  $\square$

Let us recall the definition of  $\Pi_1^1$ -additivity (see [3]): A hereditary collection  $J$  of subsets of  $X$  is called  $\Pi_1^1$ -**additive** if for every sequence  $\{A_\xi\}_{\xi < \theta}$  of sets in  $J$  such that the associated prewellordering

$$x \preceq y \text{ iff } x, y \in \bigcup_{\xi < \theta} A_\xi \text{ \& least } \xi (x \in A_\xi) \leq \text{least } \xi (y \in A_\xi)$$

is  $\Pi_1^1$ , we have that  $\bigcup_{\xi < \theta} A_\xi \in J$ . As we said before, we have the following

**Corollary 5.6.** *Let  $I$  be a  $\sigma$ -ideal of closed meager subsets of  $X$  with the covering property. Then  $I^{int}$  is  $\Pi_1^1$ -additive.*

**Proof:** The proof is the same as in the case of the  $\sigma$ -ideal of closed meager sets (see [3]). Towards a contradiction, assume  $\theta$  is the least ordinal such that there is a sequence  $\{A_\xi\}_{\xi < \theta}$  of sets in  $I^{int}$  such that the associated prewellordering  $\preceq$  is  $\Pi_1^1$ , but  $\bigcup_{\xi < \theta} A_\xi \notin I^{int}$ .

First we observe that by the same argument as in [3] we have that  $\theta$  is a limit ordinal.

Let  $K \subseteq \bigcup_{\xi < \theta} A_\xi$  with  $K \notin I$  and fix a Polish topology  $\tau$  compatible with  $I$  such that  $K$  is not  $\tau$ -meager. The restriction of  $\preceq$  to  $K \times K$  is  $\Pi_1^1$  and hence it has the property of Baire with respect to  $\tau$ . We can assume that we are working in  $(K, \tau)$ . For every  $x \in K$  we have

$$S_x = \{y \in K : y \preceq x\} \subseteq \bigcup_{\xi < \eta} A_\xi$$

for some  $\eta < \theta$  (as  $\theta$  is limit). Hence by the minimality of  $\theta$  we have that  $S_x \in I^{int}$ . From the previous theorem we get  $S_x$  is  $\tau$ -meager. By the Kuratowski-Ulam theorem (see for instance [10]) we know that for  $\tau$ -comeager many  $y$ 's,  $S^y = \{x \in K : y \preceq x\}$  is  $\tau$ -meager. So as  $K = S_y \cup S^y$ , then  $K$  is  $\tau$ -meager, which is a contradiction.  $\square$

And then we get the following

**Corollary 5.7.** *Let  $I$  be a  $\Pi_1^1$   $\sigma$ -ideal of closed meager subsets of  $X$  with the covering property. There exists a largest  $\Pi_1^1$  set in  $I^{int}$ .*

**Proof:** In order to apply theorem 1A-2 in [3] we need only to show that the collection of  $\Sigma_1^1$  sets in  $I^{int}$  is  $\Pi_1^1$  on the codes. This is a consequence of the fact that  $I$  is strongly calibrated, as we have shown it in section 2, theorem 3.2.  $\square$

When we work in  $2^\omega$ , the largest  $\Pi_1^1$  set in  $I^{int}$  can be characterized in the same fashion as  $C_1$ , the largest  $\Pi_1^1$  set without perfect subset. The main theorem is the following

**Theorem 5.8.** *Let  $I$  be a  $\Pi_1^1$   $\sigma$ -ideal of meager subsets of  $2^\omega$  with the covering property. Then there is a largest  $\Pi_1^1$  set  $C_1(I)$  in  $I^{int}$  which is characterized by*

$$x \in C_1(I) \text{ iff } \exists T \in L_{\omega_1^T} (T \text{ is a tree on } 2 \text{ \& } x \in [T] \text{ \& } [T] \in I).$$

From now on we fix a  $\Pi_1^1$   $\sigma$ -ideal  $I$  of closed meager subsets of  $2^\omega$  with the covering property.

There is a derivative operator on closed sets similar to the Cantor-Bendixson derivative which will provide us with canonical closed sets to cover a given  $\Sigma_1^1$  set in  $I^{ext}$ .

**Definition 5.9.** *Let  $S$  be a tree on  $2 \times \omega$ ; define a derivative as follows*

$$(s, u) \in S^{(1)} \text{ iff } \overline{p[S_{(s,u)}]} \notin I.$$

*By transfinite recursion we define  $S^\eta$  for every ordinal  $\eta$ .*

$$S^{\eta+1} = (S^\eta)^{(1)}.$$

*and for  $\lambda$  a limit ordinal*

$$S^\lambda = \bigcap_{\eta < \lambda} S^\eta$$

Notice that  $S^\eta$  is also a tree on  $2 \times \omega$  and  $S^{\eta+1} \subseteq S^\eta$ . Since  $S$  is countable then there is a countable ordinal  $\theta$  such that  $S^{\theta+1} = S^\theta$ . We denote this fixed point by  $S^\infty$ .

**Lemma 5.10.**  $S^\infty = \emptyset$  iff  $p[S] \in I^{ext}$ .

**Proof:** Suppose that  $S^\infty = \emptyset$ . Let  $\theta$  be a countable ordinal such that  $S^\theta = \emptyset$ . Since  $([S^\eta])$  is sequence of subsets of  $[S]$  that decreases to the empty set then we have

$$p[S] \subseteq \bigcup \{ \overline{p[S_{(s,u)}^\alpha]} : \overline{p[S_{(s,u)}^\alpha]} \in I \text{ \& } \alpha < \theta \text{ \& } (s, u) \in S \}.$$

This clearly shows that  $p[S] \in I^{ext}$ .

On the other hand suppose that  $p[S] \in I^{ext}$ . Say  $p[S] \subseteq \bigcup K_n$  with  $K_n \in I$ . Let  $L = [S^\infty]$ . We have that  $L \subseteq \bigcup (K_n \times \omega^\omega)$ . Towards a contradiction suppose that  $L \neq \emptyset$ . By the Baire category theorem there is an  $n$ ,  $(s, u) \in S^\infty$  such that  $\emptyset \neq L \cap (N_s \times N_u) \subseteq K_n \times \omega^\omega$ . Hence  $\overline{p[S_{(s,u)}^\infty]} \in I$ , which contradicts that  $(s, u) \in S^\infty$ .  $\square$

Before proving the necessary lemmas to prove theorem 5.8 let us give an idea of how the proof goes. Fix a  $\Pi_1^1$  set  $A \in I^{int}$ . Let  $T$  be a recursive tree on  $2 \times \omega$  such that

$$x \in A \text{ iff } T(x) \text{ is well-founded.}$$

Let  $x \in A$  and let  $\xi = |T(x)|$ . There is a canonical way of defining a tree  $S_\xi$  on  $2 \times \xi$  such that

$$|T(x)| \leq \xi \text{ iff } S_\xi(x) \text{ is not well-founded.}$$

Put  $S = S_\xi$ . As  $p[S]$  is a  $\Sigma_1^1$  subset of  $A$  and  $A \in I^{int}$ , then  $p[S] \in I^{ext}$ . We can easily translate the definition of the derivative to the space  $2 \times \xi$ . Hence by 5.10  $S^\infty = \emptyset$ . Thus the closed sets  $\overline{p[S_{(s,u)}^\alpha]}$  cover  $p[S]$ , as in the proof of 5.10. The key of the proof is the fact that for each of these closed sets we can find a tree  $T_{(s,u)}^\alpha$  in the least admissible set containing  $\xi$  such that

$$\overline{p[S_{(s,u)}^\alpha]} \subseteq [T_{(s,u)}^\alpha] \in I.$$

Since clearly  $\xi < \omega_1^x$ , this tree belongs to  $L_{\omega_1^x}$ , and we are done.

We will define the trees  $S_\xi$  uniformly on the codes of  $\xi$  using the following

**Lemma 5.11.** (Shoenfield, see [9]) Let  $T$  be a recursive tree on  $2 \times \omega$ . Let  $A \subseteq 2^\omega$  be defined by

$$x \in A \text{ iff } T(x) \text{ is well-founded.}$$

Define also for each countable ordinal  $\xi$

$$x \in A_\xi \text{ iff } |T(x)| \leq \xi.$$

There is a recursive relation  $S \subseteq \omega^\omega \times 2^{<\omega} \times \omega^{<\omega}$  such that

(i) if  $w \in WO$  and  $|w| = \xi$ , then  $S(w) = \{(t, s) : S(w, t, u)\}$  is a tree on  $2 \times \omega$  such that

$$x \in A_\xi \text{ iff } S(w)(x) \text{ is not well-founded.}$$

(ii) There is a tree  $S_\xi$  on  $2 \times \xi$  (as we mentioned above) such that  $p[S_\xi] = A_\xi$  and this tree belongs to the least admissible set containing  $\xi$ . Moreover, given a sequence  $u \in \omega^{<\omega}$ , by using the wellorder of  $\omega$  given by  $w$  we can think that  $u$  codes a sequence of ordinals  $h$  (and viceversa given  $h$  we can find  $u$ ) such that

$$(t, u) \in S(w) \text{ iff } (t, h) \in S_\xi.$$

Thus if  $w, z \in WO$  and  $|w| = |z| = \xi$ , then  $S(w)$  and  $S(z)$  code essentially the same tree  $S_\xi$ . □

In the following lemma we compute the complexity of the derivative defined above.

**Lemma 5.12.** *Let  $I$  be a  $\Pi_1^1$   $\sigma$ -ideal of closed subsets of  $2^\omega$  with the covering property. Let  $T$  and  $S$  as in lemma 5.11.*

(i) *There is a  $\Sigma_1^1$  relation  $P$  on  $\omega \times \omega \times \omega^\omega$  such that for  $v, w \in WO$  we have*

$$P(t, u, v, w) \text{ iff } (t, u) \in [S(w)]^{|v|}.$$

Where  $[S(w)]^{|v|}$  is defined as in 5.10.

(ii) *Let  $A$  and  $A_\xi$  be defined as in 5.11 and suppose that  $A \in I^{int}$ . For every  $\xi < \omega_1$  and every  $w \in WO$  with  $|w| = \xi$ , the closure ordinal of  $S(w)$  is  $< \xi^+$  (the least admissible ordinal bigger than  $\xi$ ).*

**Proof:** Let  $D$  be the following relation on  $\omega \times \omega \times \omega^\omega$ :

$$D(t, u, J) \text{ iff } J \text{ is a tree on } 2 \times \omega \text{ \& } (t, u) \in J^{(1)}.$$

We claim that  $D$  is  $\Sigma_1^1$ . To see this, consider the following relation

$$B(x, J) \text{ iff } J \text{ is a tree on } 2 \times \omega \text{ \& } x \in \overline{proj[J]}.$$

$B$  is clearly  $\Sigma_1^1$  and  $D(t, u, J) \text{ iff } B(J_{(t,u)}) \notin I^{int}$ . We have shown in section 2 theorem 3.2 that the collection of  $\Sigma_1^1$  sets in  $I^{int}$  is  $\Pi_1^1$  on the codes of  $\Sigma_1^1$  sets; this easily implies that  $D$  is  $\Sigma_1^1$ .

We will use the recursion theorem to define  $P$ . Let  $\mathcal{U}$  be a  $\Sigma_1^1$  universal set on  $\omega \times \omega \times \omega^\omega \times \omega^\omega \times \omega^\omega$ . Consider the following relation

$$\begin{aligned} Q(t, u, v, w, \rho) \text{ iff } & v \notin WO \text{ or } (v \in LO \text{ \& } v \equiv \emptyset \text{ \& } S(t, u, w)) \\ & \text{or } (\exists z)(v, z \in LO \text{ \& } v \equiv z + 1 \text{ \& } D(t, u, \{(l, k) : \mathcal{U}(l, k, z, w, \rho)\})) \\ & \text{or } v \text{ is limit \& } (\forall n) \mathcal{U}(t, u, v[n, w, \rho]) \end{aligned}$$

where  $v \equiv \emptyset$  means that  $v$  codes the empty order;  $v \equiv z + 1$  means that the linear order coded by  $v$  has a last element and  $z$  is the linear order obtained by deleting this last element and  $v[n]$  is the linear order obtained by restricting  $v$  to  $\{m : m <_v n\}$ .  $v$  is limit means that for all  $n$  there is  $m$  such that  $n <_v m$ .

Notice that  $D(t, u, A)$  holds iff  $\exists B(B \subseteq A \text{ \& } A \text{ is a tree \& } D(t, u, B))$  (i.e., it is a monotone operator), hence  $Q$  is  $\Sigma_1^1$ . By the recursion theorem there is a recursive  $\rho^*$  such that

$$Q(t, u, v, w, \rho^*) \longleftrightarrow \mathcal{U}(t, u, v, w, \rho^*).$$

As usual, put

$$P(t, u, v, w) \longleftrightarrow \mathcal{U}(t, u, v, w, \rho^*).$$

By induction on the length of  $v \in WO$  one can easily show that if  $w \in WO$ , then

$$P(t, u, v, w) \longleftrightarrow (t, u) \in [S(w)]^{|v|}.$$

(ii) Let  $w \in WO$  with  $|w| = \xi$  and let  $S = S(w)$ .  $A_\xi = p[S]$  is a  $\Sigma_1^1$  set in  $I^{int}$ . As  $I$  has the covering property, then by lemma 5.10  $S^\infty = \emptyset$ . Since the derivative operator is  $\Sigma_1^1$  it is a standard fact that in this case the closure ordinal of  $S$  is recursive in  $S$ , hence recursive in  $w$ .

From 5.11 we also get the following: Let  $z \in WO$  with  $|w| = |z| = \xi$  and let  $u, v \in \omega^{<\omega}$ . If  $u, v$  code the same sequence of ordinals with respect to the wellorder of  $\omega$  given by  $w$  and  $z$  respectively, then

$$(t, u) \in S(w)^{(1)} \text{ iff } (t, v) \in S(z)^{(1)}.$$

In particular the closure ordinal of  $S(w)$  and of  $S(z)$  are the same. Let then  $z$  be a generic (with respect to the partial order that collapses  $\xi$  to  $\omega$ ) ordinal code for  $\xi$ . It is a standard fact that  $\omega_1^z = \xi^+$ . This finishes the proof of (ii).  $\square$

A key fact in the proof is that the trees  $S(w)$  in the previous lemma have an invariant definition in the following sense.

**Definition 5.13.** Let  $\sim$  be an equivalence relation on  $\omega^\omega$  and  $\Gamma$  be a pointclass. We say that a set  $A$  is  $\sim$ -invariantly- $\Gamma(\alpha)$  if there is a  $\Gamma$  relation  $R$  on  $X \times \omega^\omega$  such that for every  $\beta \sim \alpha$  we have

$$x \in A \text{ iff } R(x, \beta).$$

In particular  $A$  is called  $\sim$ -invariantly- $\Delta_1^1(\alpha)$ , if  $A$  is both  $\sim$ -invariantly- $\Sigma_1^1(\alpha)$  and  $\sim$ -invariantly- $\Pi_1^1(\alpha)$ .

Consider the following equivalence relation on  $\omega^\omega$ : Let  $LO$  be the collection of codes of linear orders of  $\omega$ . We say that two codes  $\alpha$  and  $\beta$  in  $LO$  are isomorphic if the linear orders coded by them are isomorphic. Define  $\equiv$  by

$$\alpha \equiv \beta \text{ iff } \alpha, \beta \in LO \text{ \& } \alpha \text{ and } \beta \text{ are isomorphic.}$$

It is a standard fact that  $\equiv$  is a  $\Sigma_1^1$  relation (see [9]). The following two lemmas make clear why it is interesting to look at the notion of  $\equiv$ -invariantly definable sets.

**Lemma 5.14.** Let  $\xi$  be a countable ordinal and  $w$  an ordinal code for  $\xi$ . Let  $T \subseteq \omega$  be a  $\equiv$ -invariantly- $\Delta_1^1(w)$  set. Then  $T$  belongs to the least admissible set containing  $\xi$ .

**Proof:** Let  $M$  denote the least admissible set containing  $\xi$ . We will show that  $T$  is  $\Delta_1^1$  definable over  $M$ . Let  $R \subseteq \omega \times \omega^\omega$  be a  $\Pi_1^1$  set such that for every ordinal code  $w$  with  $|w| = \xi$ , we have

$$s \in T \text{ iff } R(s, w).$$

Let  $\psi$  be a  $\Sigma_1$  formula (in ZF) such that if  $N$  is an admissible set and  $w \in N$ , then

$$R(s, w) \text{ iff } N \models \psi(s, w) \quad (*)$$

Consider the notion of forcing  $\mathbf{P}$  that collapses  $\xi$  to  $\omega$ . If  $G$  is  $\mathbf{P}$ -generic, let  $w_G$  be the corresponding ordinal code, i.e.,

$$w_G(n, m) = 0 \text{ iff } \exists p \in G(p(n) < p(m)).$$

Consider the following name

$$\tau = \{\langle \sigma, p \rangle : \sigma = \langle (n, m), 0 \rangle \text{ and for some ordinals } \alpha < \beta, \langle n, \alpha \rangle, \langle m, \beta \rangle \in p\}.$$

Then for every  $\mathbf{P}$ -generic  $G$ ,  $i_G(\tau) = w_G$ . Since for every admissible set  $N$ ,  $N[G]$  is also admissible, then from (\*) we get

$$R(s, w_G) \text{ iff } M[G] \models \psi(s, w_G). \quad (**)$$

As (\*\*) holds for every  $G$   $\mathbf{P}$ -generic, then

$$s \in T \text{ iff } \Vdash \psi(\check{s}, \tau).$$

Since  $\psi$  is  $\Sigma_1$ , the relation  $B(s, \tau) \text{ iff } \Vdash \psi(\check{s}, \tau)$  is  $\Sigma_1$  over  $M$ . Hence  $T$  is  $\Sigma_1$  over  $M$ . Similarly we have that  $s \notin T$  is  $\Sigma_1$  over  $M$ . This finishes the proof.  $\square$

There is another basic fact about  $\Sigma_1^1$  equivalence relations and  $\Pi_1^1$  sets that we are going to use.

**Definition 5.15.** (Solovay [5]) Let  $\sim$  be an equivalence relation on  $\omega^\omega$  and  $P \subseteq \omega^\omega$  be a  $\sim$ -invariant set, i.e., if  $x \in P$  and  $y \sim x$  then  $y \in P$ . A norm  $\varphi : P \rightarrow \text{ordinals}$  is called  $\sim$ -**invariant** if

$$x \sim y \text{ \& } x \in P \Rightarrow \varphi(x) = \varphi(y).$$

Let  $\Gamma$  be a pointclass. We say that  $\Gamma$  is **invariantly normed** if for every equivalence relation  $\sim$  in  $\check{\Gamma}$  and every  $\sim$ -invariant set  $P$  in  $\check{\Gamma}$ ,  $P$  admits a  $\sim$ -invariant norm.

It was proved by Solovay (see [5]) that  $\Pi_1^1$  is invariantly normed.

The following result is the “invariant” version of 4.2.

**Lemma 5.16.** (see Barua-Srivatsa [1]) Let  $\sim$  be a  $\Sigma_1^1$  equivalence relation on  $\omega^\omega$  and  $T$  be a  $\sim$ -invariantly  $\Sigma_1^1(\alpha)$  binary tree with  $[T] \in I$ . There is a  $\sim$ -invariantly- $\Delta_1^1(\alpha)$  tree  $S$  such that  $[T] \subseteq [S]$  and  $[S] \in I$ .

**Proof:** The proof is entirely similar to the one of 4.2. We will sketch it, to point out where we use the notion of  $\sim$ -invariant sets.

Put a  $\sim$ -invariant  $\Pi_1^1(\alpha)$  norm over  $\text{Seq} - T$ . Define as in 4.2 the sets  $A_s$  and the set  $M$ . We claim that  $A_s$  is  $\sim$ -invariantly  $\Sigma_1^1(\alpha)$  and  $M$  is  $\sim$ -invariantly  $\Pi_1^1(\alpha)$ . Assuming the claim we finish the proof.

The separation theorem holds in an invariant form, i.e. given two disjoint  $\sim$ -invariant  $\Sigma_1^1$  sets there is a  $\sim$ -invariant  $\Delta_1^1$  set separating them. Thus, as in the proof of 4.2, let  $B$  be a  $\sim$ -invariant  $\Delta_1^1(\alpha)$  set such that  $T \subseteq B \subseteq M$ . Then let  $S$  be the tree generated by  $B$ ,  $S$  is easily seen to be  $\sim$ -invariantly  $\Delta_1^1(\alpha)$ .

It is clear that  $A_s$  is  $\sim$ -invariantly  $\Sigma_1^1(\alpha)$ , because the definition of the  $\Pi_1^1$  norm. Now for  $M$  we have

$$\begin{aligned} s \in M & \text{ iff } (\forall K)(K \subseteq [A_s] \Rightarrow K \in I) \\ & \text{ iff } (\forall K)\{\exists t(N_t \cap K \neq \emptyset) \text{ \& } t \notin A_s\} \text{ or } K \in I\} \end{aligned}$$

Now, it is clear that the relation inside curly brackets is  $\sim$ -invariantly  $\Pi_1^1(\alpha)$  (because the only thing that depends on  $\alpha$  is  $A_s$ ). This proves the claim.  $\square$

Now we are ready to give the

**Proof of theorem 5.8:** First we want to show that  $C_1(I)$  is a  $\Pi_1^1$  set in  $I^{int}$

$$x \in C_1(I) \text{ iff } \exists T \in L_{\omega^x}(T \text{ is a tree \& } x \in [T] \text{ \& } [T] \in I).$$

It is clearly  $\Pi_1^1$ , since

$$T \in L_{\omega_1^x} \text{ iff } \exists \gamma, \beta \in \Delta_1^1(x) [\gamma \in WO \ \& \ \beta \in L_{|w|} \ \& \ \beta = T].$$

Now we show that  $C_1(I) \in I^{int}$ . Put  $C = C_1(I)$ . By 5.4 it suffices to show that  $C$  is  $\tau$ -meager for every topology  $\tau$  compatible with  $I$ . Fix such a topology  $\tau$ . Define the following pre-wellordering on  $C$

$$x \leq y \text{ iff } x, y \in C \text{ and } \omega_1^x \leq \omega_1^y.$$

Since this pre-wellordering is in the  $\sigma$ -algebra generated by the  $\Sigma_1^1$  sets, it has the property of Baire with respect to  $\tau$ . Now, for every  $y \in C$

$$\{x \in C : x \leq y\} \subseteq \bigcup \{[T] : T \in L_{\omega_1^y} \ \& \ [T] \in I\}.$$

As every  $L_{\omega_1^y}$  is countable,  $\{x \in C : x \leq y\}$  is  $\tau$ -meager. Thus by the Kuratowski-Ulam theorem we have that except for a  $\tau$ -meager set of  $x$ 's  $\{y \in C : x \leq y\}$  is  $\tau$ -meager. Thus  $C$  is  $\tau$ -meager.

Finally, we need only to show that every  $\Pi_1^1$  set  $A$  in  $I^{int}$  is a subset of  $C_1(I)$ . Fix such an  $A$  and let  $T$  be a recursive tree on  $2 \times \omega$  such that

$$x \in A \text{ iff } T(x) \text{ is well-founded.}$$

Fix  $x \in A$  and let  $|T(x)| = \xi$ . Notice that  $\xi^+ < \omega_1^x$ . Let  $S$  as in 5.11, then for every ordinal code  $w$  with  $|w| = \xi$  we have that

$$A_\xi = p[S(w)].$$

As  $A_\xi \in I^{int}$  and  $I$  has the covering property, from lemma 5.10 we get that for some countable ordinal  $\theta$ ,  $S(w)^\infty = S(w)^\theta = \emptyset$ . Hence as in the proof of 5.10

$$A_\xi \subseteq \bigcup \{p[\overline{S(w)_{(s,u)}^\alpha}] : \overline{p[S(w)_{(s,u)}^\alpha]} \in I \ \& \ \alpha < \theta \ \& \ (s, u) \in S(w)\}.$$

We want to show that the sets  $[S(w)_{(s,u)}^\alpha]$  have an invariant definition in order to apply 5.16. Let  $P$  as in 5.12. Consider the following relations

$$(z_1, \dots, z_m) \equiv_w r \text{ iff } (r \in \omega^{<\omega}) \ \& \ (\forall i \leq m)(z_i \in LO \ \& \ w \in LO \ \& \ w[r(i) \equiv z_i])$$

where  $w[r(i)]$  is the initial segment of the linear order coded by  $w$  determined by  $r(i)$ , i.e.,

$$w[r(i)] = \{(l, k) : w(l, k) = w(l, r(i)) = w(k, r(i)) = 0\}.$$

Put

$$R(s, u, t, z, w, v) \text{ iff } t \in 2^{<\omega} \ \& \ lh(t) = n \ \& \ t \prec s \ \& \ (\exists r \in \omega^{<\omega})((z_1, \dots, z_n) \equiv_w r \ \& \ r \prec u \ \& \ P(s, u, v, w)).$$

Now consider the following equivalence relation on  $\omega^\omega \times \omega^\omega \times \omega^\omega$

$$(z, w, v) \sim (z', w', v') \text{ iff } z_0(0) = z'_0(0) \ \& \ (\forall 0 \leq i \leq z_0(0))(z_i, z'_i \in LO \ \& \ z_i \equiv z'_i \ \& \ w_i \equiv w'_i \ \& \ v_i \equiv v'_i).$$

Let  $(t, r) \in S(w)$  such that

$$x \in \overline{p[S(w)_{(t,r)}^{|v|}]}$$

and put

$$B = p[S(w)_{(t,r)}^{|v|}].$$

Now, by 5.11 (ii), if  $z$  codes a sequence of ordinals such that  $(z_1, \dots, z_m) \equiv_w r$ , then

$$x \in B \text{ iff } (\exists \alpha)(\forall n) R(x \upharpoonright n, \alpha \upharpoonright n, t, z, w, v).$$

Hence  $B$  is  $\sim$ -invariantly- $\Sigma_1^1$  with respect to the variables  $(z, w, v)$ . Let  $L$  be the tree of  $\overline{B}$ , clearly  $[L] \in I$ , thus by lemma 5.16 we have that there is a  $\sim$ -invariantly- $\Delta_1^1$  tree  $K$  on 2 such that  $[L] \subseteq [K]$  and  $[K] \in I$ .

By a similar argument as in the proof of lemma 5.14 we know that  $K$  belongs to the least admissible set containing all the ordinals coded by  $w, z, v$  (we need only to use the product of the notion of forcing defined in 5.14, one for each of the  $m$  ordinals coded in  $(z, w, v)$ , where  $m = lh(r) + 2$ ).

But from lemma 5.12(ii) we know these ordinals are less than  $\xi^+ < \omega_1^x$ . Therefore  $K \in L_{\omega_1^x}$ . This finishes the proof of the theorem 5.8.  $\square$

**Remark:** This proof clearly works for ideals on  $(2^\omega)^m$ .

## 6 On the strength of the covering property for $\Sigma_2^1$ sets

It is well known that the perfect set theorem for  $\Pi_1^1$  sets is equiconsistent with the existence of an inaccessible cardinal (Solovay). In fact,  $\omega_1^L < \omega_1$  iff the perfect set theorem holds for  $\Pi_1^1$  sets. In this section we will show that under the assumption that there are only countable many reals in  $L$ , any  $\Pi_1^1$   $\sigma$ -ideal of closed meager subsets of  $2^\omega$  with the covering property has also the covering property for  $\Sigma_2^1$  sets. Also, we will see that for some  $\sigma$ -ideals, the covering property for  $\Pi_1^1$  sets fails in  $L$  and thus it can not be proved in ZFC.

**Theorem 6.1.** *Let  $I$  be a  $\Pi_1^1$   $\sigma$ -ideal of meager closed subsets of  $2^\omega$  with the covering property. If  $\omega_1^L < \omega_1$ , then  $I$  has the covering property for  $\Pi_1^1$  sets. And by relativization, given  $x \in \omega^\omega$ , if  $\omega_1^{L(x)} < \omega_1$ , then the covering property holds for  $\Pi_1^1(x)$  sets.*

*Also the same result holds for  $\sigma$ -ideals of closed meager subsets of  $(2^\omega)^m$ .*

**Proof:** It clearly suffices to show that the largest  $\Pi_1^1$  set  $C_1(I)$  in  $I^{int}$  belongs to  $I^{ext}$ . But if  $\omega_1^L < \omega_1$ , then there are only countable many binary trees in  $L$ . Hence from theorem 5.8 we easily get that  $C_1(I) \in I^{ext}$ .  $\square$

The next result is a generalization of the result of Solovay that says that if there are only countable reals in  $L$ , then  $\omega^\omega \cap L$  is the largest countable  $\Sigma_2^1$  set. A similar result holds for some  $\sigma$ -ideals defined by games (see [4]).

**Theorem 6.2.** *Under the hypothesis of 6.1 the largest  $\Sigma_2^1$  in  $I^{ext}$  and in  $I^{int}$  is*

$$C_2(I) = \{x \in 2^\omega : \exists T \in L (T \text{ is a tree on } 2 \text{ \& } x \in [T] \text{ \& } [T] \in I)\}.$$

In particular, the covering property holds for  $\Sigma_2^1$  sets. And by relativization, given  $x \in \omega^\omega$ , if  $\omega_1^{L(x)} < \omega_1$ , then the covering property holds for  $\Sigma_2^1(x)$  sets.

**Proof:** If there are only countable many reals in  $L$ , then there are only countable many binary trees in  $L$ . Thus  $C_2(I)$  is clearly a  $\Sigma_2^1$  set in  $I^{ext}$ .

Let  $A$  be a  $\Sigma_2^1$  set in  $I^{int}$  and let  $B \subseteq X \times 2^\omega$  be a  $\Pi_1^1$  set such that  $x \in A$  iff  $\exists \alpha(x, \alpha) \in B$ . Let  $J$  be the  $\sigma$ -ideal of closed subsets of  $2^\omega \times 2^\omega$  defined by

$$K \in J \text{ iff } proj(K) \in I. \quad (*)$$

By proposition 3.8  $J$  has the covering property and clearly  $J$  is a  $\Pi_1^1$   $\sigma$ -ideal of meager sets. Hence by the previous theorem  $J$  has the covering property for  $\Pi_1^1$  sets. As  $A \in I^{int}$ , then  $B \in J^{int}$ . Let  $C_1(J)$  be the largest  $\Pi_1^1$  set in  $J^{int}$ , i.e.,

$$C_1(J) = \{(x, \alpha) : \exists S \in L_{\omega_1^{(x, \alpha)}}(S \text{ is a tree on } 2 \times 2 \text{ \& } (x, \alpha) \in [S] \text{ \& } proj([S]) \in I)\}.$$

It is clear that  $A \subseteq proj(C_1(J))$ . Now, let  $K$  be a closed subset of  $2^\omega \times 2^\omega$  and let  $S$  be the tree of  $K$ . Put  $T = \{t : \exists s(t, s) \in S\}$ .  $T$  is clearly a tree and by using Konig's lemma it is easy to check that  $[T] = proj([S])$ . Clearly if  $S \in L$ , then so does  $T$ . Hence

$$A \subseteq proj(C_1(J)) \subseteq \{x \in 2^\omega : \exists T \in L (x \in [T] \text{ \& } [T] \in I)\}.$$

□

The next lemma will be used in the proof that for some ideals the covering property for  $\Pi_1^1$  set fails in  $L$ . These results are due to Dougherty and Kechris, we include its proof with theirs permission.

Let us denote by  $\leq_T$  the relation of Turing reducibility, i.e.,  $x \leq_T y$  iff  $x$  is recursive in  $y$ .

**Lemma 6.3.** (Dougherty, Kechris) Let  $\mu$  be the product probability measure on  $2^\omega$  and let  $I$  be the  $\sigma$ -ideal of closed  $\mu$ -measure zero subsets of  $2^\omega$ . Then for every  $x \in 2^\omega$ ,  $\{y : x \leq_T y\} \notin I^{ext}$ .

**Proof:** Let  $\{K_n\}$  be a countable collection of sets in  $I$ . We will define  $y \notin \bigcup_n K_n$  such that  $x \leq_T y$ .

By the  $n$ -th block we mean the interval  $[2^n, 2^{n+1})$ . Call  $z \in 2^\omega$  **good** if for infinite many  $n$ 's,  $z$  is constant in the  $n$ -th block. If  $z$  is good, let  $\tilde{z}$  be defined as follows : Let  $n_0 < n_1 < \dots$  be an enumeration of the blocks on which  $z$  is constant; put  $\tilde{z}(i) = j$  if  $z$  is constantly equal to  $j$  in the  $n_i$ -th block.

We will define by induction a good  $y \notin \bigcup_n K_n$  such that  $\tilde{y} = x$ . Clearly  $x \leq_T y$  and we will be done. For every  $n$  and  $k$  with  $k > n$  and every sequence  $s \in 2^{2^n}$ , let

$$F_k^s = \{z \in 2^\omega : z \text{ is not constant in the } j\text{-th block for } n \leq j \leq k \text{ \& } s \prec z\}.$$

There are exactly  $2^{2^n} - 2$  non constant sequences of length  $2^n$ . Therefore, if  $z \in F_n^s$ , then  $z$  can take  $2^{2^j} - 2$  possible values in the  $j$ -th block. From this, one easily gets that

$$\mu(F_k^s) = (2^{2^n} - 2)(2^{2^{n+1}} - 2) \cdots (2^{2^k} - 2)/2^{2^{k+1}}.$$

Hence

$$\mu(F_k^s) = \frac{1}{2^{2^n}} \prod_{j=n}^k (1 - \frac{2}{2^{2^j}}). \quad (*)$$

If  $k \rightarrow \infty$ , the infinite product (\*) is equiconvergent with

$$\sum_{j=n}^{\infty} \frac{1}{2^{2^j}}.$$

Hence, for every  $s \in 2^n$  we have

$$\mu\left(\bigcap_{k=n}^{\infty} F_k^s\right) > 0.$$

Let  $F_s = \bigcap_{k=n}^{\infty} F_k^s$ . Now we start defining  $y$ . As  $\mu(F_{\emptyset}) > 0$ , there is  $z \in F_{\emptyset} - K_0$ . Choose  $n_0$  large enough such that if  $z \upharpoonright 2^{n_0} \prec w$ , then  $w \notin K_0$ . Define  $t_0 \in 2^{n_0+1}$  by  $t_0 \upharpoonright 2^{n_0} = z \upharpoonright 2^{n_0}$  and  $t(i) = x(0)$  for every  $i \in [2^{n_0}, 2^{n_0+1})$ . Put  $y \upharpoonright 2^{n_0+1} = t_0$ . Notice that  $t_0$  is not constant in any  $j$ -block for  $j < n_0$ . Clearly we can repeat this for  $K_1$  and  $F_{t_0}$ . So let  $z \in F_{t_0} - K_1$  and  $n_1 > n_0 + 1$  large enough such that if  $z \upharpoonright 2^{n_1} \prec w$ , then  $w \notin K_1$ . Define as before  $t_1 \in 2^{n_1+1}$  by  $t_1 \upharpoonright 2^{n_1} = z \upharpoonright 2^{n_1}$  and  $t_1(i) = x(1)$  for every  $i \in [2^{n_1}, 2^{n_1+1})$ . Put  $y \upharpoonright 2^{n_1+1} = t_1$ . The induction step should be now clear. So we get  $y \notin \bigcup_n K_n$  and  $\tilde{y} = x$ . This finishes the proof.  $\square$

As we have mentioned before, for the  $\sigma$ -ideal of countable closed subsets of  $2^\omega$  the largest  $\Pi_1^1$  set without perfect subset is characterized by

$$C_1 = \{\alpha \in 2^\omega : \alpha \in L_{\omega_1^\alpha}\}.$$

The next theorem shows that (in L)  $C_1$  cannot be covered by countable many closed sets of (Lebesgue) measure zero. However, let us observe that as  $C_1$  has no perfect subsets, it clearly has measure zero and also belongs to  $I^{int}$  for every ideal containing all singletons.

**Theorem 6.4.** (*Dougherty, Kechris*) *Let  $\mu$  and  $I$  as in 6.3. In L,  $C_1 \notin I^{ext}$ . Therefore, if  $J$  is a  $\sigma$ -ideal on  $2^\omega$  such that  $J$  contains all singletons and  $J \subseteq I$ , then (in L)  $J$  does not have the covering property for  $\Pi_1^1$  sets.*

**Proof:** Let  $\{K_n\}$  be a countable collection of closed sets of  $\mu$ -measure zero. We will show that there is  $y \in C_1$  and  $y \notin \bigcup_n K_n$ .

Let  $\{T_n\}$  be the corresponding trees and let  $\alpha < \omega_1^L$  be an ordinal such that each  $T_n \in L_\alpha$ . We can assume without loss of generality that  $\alpha$  is an index (i.e., there is  $x \in \omega^\omega$  such that  $x \in L_{\alpha+1} - L_\alpha$ ). Let  $x$  be a complete set of index  $\alpha$  (that is:  $x \in L_{\alpha+1} - L_\alpha$  and every  $y \in \omega^\omega \cap L_{\alpha+1}$  is arithmetical in  $x$ ), in particular  $\alpha < \omega_1^x$ .

Let  $y$  be as in the proof of the previous proposition. It is easy to check that  $y$  can be found in  $L_{\alpha+\omega}$ . As  $\omega_1^x \leq \omega_1^y$  (because  $x \leq_T y$ ), then  $\alpha + \omega \leq \omega_1^y$ . Hence  $y \in L_{\omega_1^y}$ , so  $y \in C_1$ . By construction  $y \notin \bigcup_n K_n$ .  $\square$

These theorems can be easily transferred to compact intervals of the real line as follows: Say we are working on  $[0, 1]$  and consider the function  $f : 2^\omega \rightarrow [0, 1]$  defined by

$$f(\varepsilon) = \sum_{i=0}^{\infty} \varepsilon(i) 2^{-(i+1)};$$

$f$  is continuous and surjective. Now, given a  $\sigma$ -ideal  $I$  of closed meager subsets of  $[0, 1]$ , define an ideal  $J$  of closed subsets of  $2^\omega$ , as follows:

$$K \in J \text{ iff } f[K] \in I.$$

Observe that  $J$  consists of meager sets (because for every nbhd  $N_s$  on  $2^\omega$  we have that  $f[N_s]$  contains an interval).

**Lemma 6.5.** *If  $I$  has the covering property, then so does  $J$ .*

**Proof:** First we show that if  $A$  is a  $\Sigma_1^1$  set, then  $A \in J^{int}$  iff  $f[A] \in I^{int}$ . The direction  $\Leftarrow$  is obvious by the definition of  $J$ .

Let  $A$  be a  $\Sigma_1^1$  set such that  $f[A] \notin I^{int}$ , say  $K \subseteq f[A]$  is a closed set and  $K \notin I$ . Define  $R$  as follows:

$$R(x, \alpha) \text{ iff } \alpha \in A \text{ \& } x \in K \text{ \& } f(\alpha) = x.$$

Then  $x \in K$  iff  $\exists \alpha R(x, \alpha)$ . Hence, as  $I$  is strongly calibrated, there is a closed set  $F \subseteq R$  such that

$$K_0 = \{x : \exists \alpha (x, \alpha) \in F\} \notin I$$

Notice that  $K_0 \subseteq K$ . Put  $L = \{\alpha : \exists x (x, \alpha) \in F\}$ . Then  $f[L] = K_0$  and  $L \subseteq A$ , so  $A \notin J^{int}$ .

The covering property for  $J$  now follows: If  $A \in J^{int}$  is a  $\Sigma_1^1$  set, then  $f[A] \in I^{int}$ . Hence  $f[A] \in I^{ext}$ , which clearly implies that  $A \in J^{ext}$ . □

**Theorem 6.6.** *Let  $I$  be a  $\Pi_1^1$   $\sigma$ -ideal of closed meager subsets of  $[0, 1]$  with the covering property. Let  $f$  be the function defined above. The largest  $\Pi_1^1$  set in  $I^{int}$  is*

$$C_1(I) = \{x \in [0, 1] : \exists T \in L_{\omega_1^x} (T \text{ is a tree on } 2 \text{ \& } x \in f[T] \text{ \& } f[T] \in I)\}$$

and the largest  $\Sigma_2^1 \in I^{ext}$  is characterized by

$$C_2(I) = \{x \in [0, 1] : \exists T \in L (T \text{ is a tree on } 2 \text{ \& } x \in f[T] \text{ \& } f[T] \in I)\}.$$

In particular, if  $\omega_1^L < \omega_1$ , then  $I$  has the covering property for  $\Sigma_2^1$  sets. And by relativization, given  $x \in \omega^\omega$ , if  $\omega_1^{L(x)} < \omega_1$ , then the covering property holds for  $\Sigma_2^1(x)$  sets.

**Proof:** First, as in the proof of theorem 5.8 we have that  $C_1(I)$  is a  $\Pi_1^1$  set in  $I^{int}$ . To see that it is the largest, consider the  $\sigma$ -ideal  $J$  defined on  $2^\omega$  as in 6.5.  $J$  has the covering property. Let  $C_1(J)$  be the largest  $\Pi_1^1$  set in  $J^{int}$  given by theorem 5.8 i.e.,

$$C_1(J) = \{\alpha \in 2^\omega : \exists T \in L_{\omega_1^\alpha} (T \text{ is a tree on } 2 \text{ \& } \alpha \in [T] \text{ \& } [T] \in J)\}.$$

Let  $A$  be a  $\Pi_1^1$  set in  $I^{int}$ . Put  $B = f^{-1}(A)$ ,  $B$  is a  $\Pi_1^1$  set in  $J^{int}$ . So  $B \subseteq C_1(J)$ , hence it suffices to show that  $f(C_1(J)) \subseteq C_1(I)$ . Let  $\alpha \in C_1(J)$  and let  $T \in L_{\omega_1^\alpha}$  such that  $\alpha \in [T]$  and  $[T] \in J$ . As  $f$  is  $\Delta_1^1$ , then  $\omega_1^\alpha = \omega_1^{f(\alpha)}$ . So  $T \in L_{\omega_1^{f(\alpha)}}$ . Thus  $f(\alpha) \in f[T]$  and also  $f[T] \in I$ .

The proof for  $C_2(I)$  is similar. □

Theorem 6.4 can also be transferred to  $[0, 1]$  as follows: Let us observe that for every basic nbhd  $N_s$  in  $2^\omega$  we have that  $\mu(N_s) = \lambda(f[N_s])$ , where  $\mu$  is the standard product measure on  $2^\omega$  and  $\lambda$  is the Lebesgue measure on  $[0, 1]$ . One easily checks that if  $f[C_1]$  can be covered by countably many closed sets of Lebesgue measure zero, then  $C_1$  can also be covered by countably many closed sets of  $\mu$ -measure zero. It is also clear that this set does not contain a perfect subset. We collect these facts in the following

**Theorem 6.7.** *Let  $I$  be a  $\sigma$ -ideal of closed subsets of  $[0,1]$  such that every set in  $I$  has Lebesgue measure zero. In  $L$ ,  $I$  does not have the covering property for  $\Pi_1^1$  sets.*

□

**Remark:** As we have already mentioned the  $\sigma$ -ideal of closed set of extended uniqueness has the covering property (see [2]). Hence, from 6.6 and 6.7 we get that the covering property for  $\Pi_1^1$  sets of extended uniqueness is not provable in ZFC, but can be proved from the hypothesis that there are only countably many reals in  $L$ . Also we get a characterization of the largest  $\Pi_1^1$  set of extended uniqueness as in 6.6.

## References

- [1] R. Barua and V.V. Srivatsa, Definable hereditary families in the projective hierarchy, preprint.
- [2] G. Debs and J. Saint Raymond. Ensembles d'unicité et d'unicité au sens large, Ann. Inst. Fourier, Grenoble, 37(3), 1987, 217-239.
- [3] A. S. Kechris, The theory of countable analytical sets, Trans. Amer. Math. Soc., 202, (1975), 259-297.
- [4] A. S. Kechris, On a notion of smallness for subsets of the Baire space, Trans. Amer. Math. Soc., 229, (1977), 191-207.
- [5] A. S. Kechris, Countable ordinals and the analytical hierarchy(II), Annals of Mathematical Logic, 15, (1978), 193-223.
- [6] A. S. Kechris, The descriptive set theory of  $\sigma$ -ideals of compact sets, Logic Colloquium'88, Ferro, Bonotto, Valentini and Zanardo (Editors), North-Holland, 1979, 117-138.
- [7] A. S. Kechris and A. Louveau, Descriptive set theory and the structure of sets of uniqueness, LMS Lecture Notes, 128, 1987.
- [8] A. S. Kechris, A. Louveau and W. H. Woodin, The structure of  $\sigma$ -ideals of compact sets, Trans. Amer. Math. Soc., 301(1), (1987), 263-288.
- [9] Y. N. Moschovakis, Descriptive set theory, North Holland, Amsterdam, 1980.
- [10] J. Oxtoby, Measure and category, Springer-Verlag, New York, 1971.